

1

2

3

CRCM5484: A BET- BDII Selective Compound

4

With Differential Anti-Leukemic Drug Modulation

5

6 *Kendall Carrasco¹; Camille Montersino^{1,2}; Carine Derviaux^{1,2}; Magali Saez-Ayala¹; Laurent Hoffer¹; Audrey*

7

Restouin^{1,2}; Rémy Castellano^{1,2}; Justine Casassa¹; Philippe Roche¹; Eddy Pasquier¹; Sébastien Combes¹;

8

Xavier Morelli^{1,2}; Yves Collette^{1,2*} & Stéphane Betzi^{1*}*

9

¹CNRS, INSERM, Aix-Marseille Univ, Institut Paoli-Calmettes, CRCM, 13009 Marseille, France

10

²Institut Paoli-Calmettes, plateforms HiTS & TrGET, 13009 Marseille, France

11

- 1 KEYWORDS:
- 2 Bromodomain, protein-protein interaction, hit-to-lead, optimization, focused library, diversity-oriented
- 3 library, drug discovery, BDII selective inhibitor, modulation, potentiation, bone-marrow-derived stromal
- 4 cell conditioned medium.
- 5

1 ABSTRACT

2 Differentially screening the Fr-PPIChem chemical library on the BET BRD4-BDII versus -BDI
3 bromodomains led to the discovery of a BDII selective tetrahydropyridothienopyrimidinone (THPTP)-
4 based compound. Structure-activity relationship (SAR) and hit-to-lead approaches allowed us to develop
5 CRCM5484, a potent inhibitor of BET proteins with a preferential and 475-fold selectivity for the second
6 bromodomain of the BRD3 protein (BRD3-BDII) over its first bromodomain (BRD3-BDI). Its very low
7 activity was demonstrated in various cell-based assays, corresponding with recent data describing other
8 selective BDII compounds. However, screening on a drug sensitivity and resistance-profiling platform
9 revealed its ability to modulate the anti-leukemic activity in combination with various FDA-approved
10 and/or in-development drugs in a cell- and context-dependent differential manner. Altogether, the
11 results confirm the originality of the THPTP molecular mode of action in the BD cavity and its potential
12 as starting scaffold for the development of potent and selective bromodomain inhibitors.

13

1 INTRODUCTION


2 The bromodomain and extra-terminal (BET) family is composed of four epigenetic reader proteins,
3 ubiquitous BRD4, BRD3 and BRD2 and testis-restricted BRDT. Each protein of this family includes a
4 tandem small protein domain called bromodomain (BD) labeled BDI and BDII, both localized at the N-
5 terminus of the proteins, and this domain can recognize and bind acetylated lysine on the histone tails.
6 When BDs are bound to chromatin, they recruit transcription factors and play a role in gene
7 transcription¹. Pan-BET inhibitors are small compounds that can bind the eight BET BDs with similar
8 potencies. This class of BD inhibitors has already been extensively studied for their therapeutic potential
9 in both inflammation²⁻⁸ and oncology applications⁹⁻¹⁴. Among these well-studied compounds are
10 JQ1^{10,15}, OTX-015¹⁶ and I-BET-151¹⁷. Some of these molecules are currently being evaluated in clinical
11 trials with oncology indications, confirming the interest for this protein family as therapeutic targets¹⁸⁻²⁰.
12 Nevertheless, toxicities and resistance phenomena have also been reported^{19,21-23}. One of the main
13 strategies to understand and overcome these issues has been to elucidate the functional contribution of
14 the individual BDI and BDII domains in the BET family with the purpose of improving the adverse event
15 profile observed in clinical trials.

16 Despite a high homology across the BDs in the BET family, several selective inhibitors of the BDI or
17 BDII BET domains have recently emerged in the literature²⁴⁻²⁷, but few are currently engaged in clinical
18 trials. The first developed selective compounds were BDI selective, the most extensively studied are
19 olinone²⁸, MS402²⁹, and GSK778³⁰. Developing BDII selective compounds was far more challenging, and
20 only a handful of BDII selective inhibitors have been developed to date (Figure 1). The oldest compound,
21 RVX-208 based on a quinazolinone chemical core, exhibited a selectivity of 20-fold with K_D values of
22 4100 nM and 200 nM (as measured by microcalorimetry (ITC)) against BRD4-BDI and BRD4-BDII,
23 respectively^{14,31}. GSK340 with a tetrahydroquinoxaline core is another BDII selective compound that
24 exhibited K_D values of 3000 nM and 63 nM against BRD4-BDI and BRD4-BDII, respectively³² (as
25 determined by BROMOScan). Very recently, ABBV-744 (tetrahydropyrrolopyridinone) was reported as a

1 nanomolar BDII selective inhibitor and clinical candidate with a K_D measured by TR-FRET at 520 nM and
2 1.6 nM against BRD4-BDI and BRD4-BDII, respectively^{33,34}. Finally, GSK046 (fluorobenzamide) and
3 GSK620 (pyridinone), the latest additions to this list of BDII selective compounds, displayed IC_{50} values
4 against the BDs of BRD3 that were 3600 nM and 98 nM for GSK046, and 2500 nM and 72 nM for
5 GSK620, as determined by TR-FRET^{30,35}. These compounds were evaluated for application in
6 inflammatory disease, and the authors concluded that inhibition of the BET BDII domain may be a useful
7 therapeutic strategy to counter inflammatory damage³⁰.

8 In order to seek for new potential cores selective of either BDI or BDII domains, we miniaturized and
9 robotized a 1536 BD inhibitors assay and initiated a differential (BDI vs BDII) high-throughput screen
10 (HTS) of the Fr-PPICChem library. Fr-PPICChem is a unique chemical library composed of 10314 diverse
11 small molecules dedicated to protein-protein interaction (PPI) inhibition that displays considerably
12 higher hit rates against PPIs than that of classical libraries³⁶. Another fundamental advantage of this
13 library is that its compounds as well as a series of structural analogs are commercially available.
14 Moreover, medicinal chemistry filters were applied to remove undesirable structures (such as Pan-Assay
15 Interference Compounds [PAINS], frequent hitters and toxic compounds) and to improve drug-likeness
16 properties. From the Fr-PPICChem screen, an original tetrahydropyridothienopyrimidinone (THPTP)-based
17 compound was identified as a bromodomain binder with a BDII selective profile that was further
18 optimized through a structure activity relationship (SAR) campaign assisted by X-ray structural data. Our
19 optimized hit - CRCM5484 - exhibits similar *in vitro* and in cellular activity to the most selective
20 compounds described in the literature. Finally, a drug sensitivity response profile (DSRP) in a
21 combination screen of CRCM5484 with 78 pharmacological agents pinpointed that CRCM5484 could
22 potentiate the antiproliferative activity of certain drugs in patient-derived leukemia cells, which could be
23 exploited for future developments. This potentiating effect is dependent upon the cellular context, and
24 the drug response could be further potentiated with the addition of culture medium conditioned by

- 1 human bone marrow stromal cells to mime the microenvironment of tumor cells; this highlights the
- 2 importance of considering the impact of the tumor microenvironment when evaluating drug sensitivity
- 3 in oncology.



- 4
- 5 **Figure 1. Chemical structures of the literature BDII selective inhibitors RVX-208, GSK340, ABBV-744, GSK046 and GSK620.**
- 6 Chemical moieties mimicking the histone acetylated lysine interaction with the key asparagine amino acid of the BDs are
- 7 highlighted in orange.


8

1 RESULTS

2 **THPTP core was identified by a “Fr-PPIChem” screening as a candidate for selective BDII BET**
3 **inhibition**

4 To develop our BDII selective BET inhibitor, we performed a differential screening on the “Fr-
5 PPIChem” chemical library³⁶. We used homogeneous time-resolved fluorescence (HTRF)³⁷ to screen the
6 10314 compound library against both BDs of BRD4 (BRD4-BDI and BRD4-BDII). The scatter plot in Figure
7 2A confirms that the hits in the eight 1536-well screening plates were randomly distributed, thus
8 experimental bias could be excluded. In addition, we measured several statistical data quality features
9 for the screening. All these indicators were in agreement with industry standards, including Z' factors of
10 0.83 and 0.87 and signal-to-noise ratios of 12.5 and 7.8 for BRD4-BDI and BRD4-BDII, respectively. The
11 variability coefficient was also found to be below 5% for the whole screening. The screen designed to
12 highlight BDII selective compounds identified 17 molecules with >30% BRD4-BDII inhibition and <30%
13 BRD4-BDI screening at a final concentration of 1.25 μ M. Among the molecules that met these criteria, 7
14 were selected for further evaluation (Table 1 and Figure S1 and S2A). The same selection threshold used
15 to select BDI selective compounds would have led to 132 molecules, highlighting the challenge of
16 developing BDII over BDI selective inhibitors.

17 We focused on compound **1** (Figure 2B) because it had a high affinity for BRD4-BDII (~900 nM) and
18 because of its original THPTP chemical scaffold in the BET inhibitor chemical landscape. Indeed, only one
19 related compound could be identified for this chemical scaffold in the literature as a BD binder in one
20 crystal structure deposited in the PDB in 2016 (PDB ID: 4ZW1), but there was no follow-up study or
21 publication exploiting these data (Figure S3A). The other identified compounds, a dimethyl isoxazole
22 core of **2**, **3** and **4**, has already been described in the BD inhibitor I-BET-151²⁵ or GSK778³⁰, and the
23 trimethoxyphenyl core of **5** is found in DC-BD-03³⁸ (Figure S2B). These results demonstrate the potential
24 of the Fr-PPIChem chemical library to identify a new chemical core during a screening campaign.



1
2 **Figure 2. Fr-PPICChem screening and compound 1 characterization.** (A) Scatter plot of Fr-PPICChem screening results against
3 BRD4-BDI (empty blue circles) and BRD4-BDII (empty green circles). BDII selective compounds identified above the 30%
4 threshold are shown as solid green circles. (B) Compound 1 chemical structure. (C) HTRF selectivity assay on the BDI (solid lines)
5 and BDII (dashed lines) of BRD2 (blue), BRD3 (green) and BRD4 (red). (D) BROMOscan data on the BET bromodomain family for
6 both BDs of BRD2, BRD3 and BRD4.

7 **Table 1. Screening results of seven BDII selective hits identified.** The IC_{50}
8 were obtained by HTRF assay on BRD4-BDI and BRD4-BDII.

Compound	IC_{50} (μM)		BDII selectivity (fold)
	BRD4-BDI	BRD4-BDII	
1	4.0 \pm 0.5	0.9 \pm 0.1	4.4
2	11.0 \pm 0.9	0.9 \pm 0.1	12.0
3	9.3 \pm 0.8	1.1 \pm 0.1	7.8
4	7.4 \pm 0.5	1.5 \pm 0.1	4.9
5	4.3 \pm 0.2	0.8 \pm 0.1	5.4
6	8.2 \pm 0.8	0.9 \pm 0.1	9.0
7	1.1 \pm 0.1	3.2 \pm 0.5	0.4

9
10 The affinity and the selectivity of **1** were confirmed by two experiments: an HTRF selectivity assay on
11 the six BDs of the BET family and a BROMOscan assay on the whole BD family tree (DiscoverX - Eurofins)
12 (Figure 2C and D). These results confirm that **1** preferentially binds the BDII domains of BRD3 and BRD2,

1 as confirmed by the IC₅₀ values of 43 nM and 120 nM, respectively, which were measured by HTRF. The
2 affinity for BRD4-BDII is significantly lower, with an IC₅₀ of 930 nM. This was further supported by the
3 BROMOscan results, in which the dissociation constants for the BDIIs of BRD4, BRD3 and BRD2 were
4 measured at 2600 nM, 325 nM and 775 nM, respectively. This characteristic is common to other
5 already-published BDII selective compounds, such as GSK046 and GSK620 that also bind preferentially
6 the BDII sub-domains of BRD2 and BRD3 over BRD4-BDII³⁰.


7 **THPTP optimization by SAR**

8 To investigate the molecular mode of action of **1** and how its different chemical functions affect the
9 affinity and selectivity toward BET BDs, we initiated an SAR campaign by ordering similar compounds
10 from commercial libraries (SAR by catalog). The first five compounds purchased were selected because
11 of the common 7-alcanoyl-5,6,7,8-tetrahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-4(1H)-one core, as
12 shown in Table 2. After measuring their affinities for both BDs of BRD3 by HTRF, we concluded that the
13 *N*-acetylated pyrido moiety is a key binding component. This observation suggests there is a different
14 binding mode than that of the 4ZW1 PDB compound in which the interaction is driven by the xlenol
15 core (Figure S3).

16 The presence of a furfuryl group as an R substituent also appears to be important for the selectivity
17 profile. This substitution is indeed the only difference from **1**, and the affinity for BRD3-BDII is degraded
18 with IC₅₀ values of 85 nM, 410 nM and 6300 nM for **1**, **8** and **9**, respectively. Compound **12** is the best
19 compound in this series, exhibiting an IC₅₀ of 140 nM on BRD3-BDII and a selectivity of 100-fold. We
20 hypothesized that the furan moiety adopts a position in close proximity to histidine H433 of BRD2-BDII,
21 one of the main hot-spot amino acids that is targeted when developing BDII selective drugs. H433 is
22 indeed one of the few residues discriminating BDII from BDI in an otherwise identical cavity. The only
23 difference with **1** is the unsubstituted phenyl ring (Table 2). This observation suggests that aryl is

1 important for binding to BD. The HTRF assay on BRD4, BRD3 and BRD2 for both BDs shows that **12**
 2 maintains the BDII selective profile (Figure S4A).


3 **Table 2. Structure Activity Relationship (SAR) by catalog.** The IC₅₀ were obtained by HTRF assay on BRD3-BDI and BRD3-BDII.



Compound	R	Ar	IC ₅₀ (nM)		BDII selectivity (fold)
			BRD3-BDI	BRD3-BDII	
1	2-furfuryl	5-chloro-2-methylphenyl	8500 ± 130	85 ± 3	100
8	phenyl	5-chloro-2-methylphenyl	> 25000 ± 1487	6300 ± 220	> 4
9	tetrahydrofuran-2-ylmethyl	5-chloro-2-methylphenyl	7800 ± 400	410 ± 10	19
10	cyclohexyl	-	> 25000 ± 2435	NA	NA
11	phenyl	phenyl	> 25000 ± 1726	5500 ± 3638	> 5
12	2-furfuryl	phenyl	14000 ± 510	140 ± 10	100

5 The importance of aryl prompted us to further investigate with another round of SAR by catalog,
 6 focusing on various phenyl substitutions. As the HTRF evaluation of the previous series identified
 7 solubility issues, we also included the predicted compound log P as an additional selection metric. We
 8 chose compounds with a log P below 4 because the predicted log P of **1** is 4.2 (Table3). The affinity and
 9 selectivity evaluation of this series by HTRF provided valuable SAR information. First, methyl, ethoxy or
 10 methoxy substituents in the *-Para* position degraded the affinity for BRD3-BDII, with IC₅₀ values ranging
 11 from 570 to 880 nM against the 190 nM measured for **12**. This suggests the ring has a close proximity to
 12 the protein surface, causing steric hindrance when substituents are included. Substitution with a
 13 methoxy or ethoxy groups in the *-Ortho* position could increase the compound affinity because of
 14 additional hydrogen bonds. Compound **16**, with anisole, is the most interesting compound of this set,
 15 with affinities of 2 200 nM and 72 nM for BRD3-BDI and BRD3-BDII, respectively. The HTRF assay on both
 16 BDs of the BET family validated the BDII selective profile of **16** (Figure S4B).


1 **Table 3. Structure Activity Relationship (SAR) of compound 12 benzyl ring.** IC₅₀ values were obtained by HTRF on BRD3-BDI
 2 and BRD3-BDII. Log P values were generated with Marvin Sketch. Compounds are sorted according to their affinity for BRD3-
 3 BDII.



Compound	Ar	Log P	IC ₅₀ (nM)		BDII selectivity (fold)
			BRD3-BDI	BRD3-BDII	
1	5-Chloro-2-methylphenyl	4.2	8500 ± 130	85 ± 3	100
12	phenyl	3.0	2100 ± 230	190 ± 12	11
13	3-Methoxyphenyl	2.9	2800 ± 330	300 ± 12	9
14	2-Chlorophenyl	3.7	2300 ± 180	100 ± 5	23
15	3-Methylphenyl	3.6	5200 ± 2100	440 ± 34	12
16	2-Methoxyphenyl	2.9	2200 ± 150	72 ± 3	31
17	4-Methoxyphenyl	2.9	3600 ± 910	570 ± 22	6
18	4-Fluorophenyl	3.2	3000 ± 2300	420 ± 18	7
19	2-Methylphenyl	3.6	3200 ± 750	220 ± 7	15
20	3-Methylphenyl	3.6	3100 ± 970	630 ± 23	5
21	3-Chlorophenyl	3.7	4400 ± 1700	490 ± 15	9
22	2-Ethoxyphenyl	3.2	3500 ± 730	150 ± 9	23
23	2-4-Difluorophenyl	3.3	7500 ± 3300	380 ± 13	13
24	2-Fluorophenyl	3.2	4200 ± 320	210 ± 6	20
25	3-Fluoro-4-methylphenyl	3.7	6500 ± 5200	210 ± 6	31
26	3-Methylthiophenyl	3.7	4500 ± 1900	410 ± 12	11
27	4-Chlorophenyl	3.7	4500 ± 2000	530 ± 18	8
28	2-5-Difluorophenyl	3.3	4100 ± 1600	170 ± 4	24
29	3-Acetylphenyl	2.6	1500 ± 64	170 ± 6	9
30	4-Acetylphenyl	2.6	1300 ± 45	320 ± 12	4
31	4-Ethoxyphenyl	3.2	3800 ± 2500	880 ± 22	4
32	3-Chloro-4-fluorophenyl	3.8	4900 ± 2700	740 ± 28	7
33	3,4-Difluorophenyl	3.3	3800 ± 1500	470 ± 11	8

7
8 At this point, all purchasable analogs were evaluated, and we expanded the SAR study with in-house
 9 chemical synthesis. We pursued the optimization strategy by exploring additional modifications of the
 10 aromatic ring to increase the activity and solubility. With the addition of the piperidine group, the
 11 predicted Log P decreased from 2.9 for compound 16 to 2.4 for compound 34 (CRCM5484). The 2-((7-

1 acetyl-3-(furan-2-ylmethyl)-4-oxo-3,4,5,6,7,8-hexahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-2-
2 yl)thio)-*N*-(2-methylpyridin-3-yl)acetamide (CRCM5484, **34**) was prepared by convergent synthesis in
3 three steps as depicted in scheme 1. A mixture of commercially available *N*-acetyl-4-piperidone,
4 activated ethyl 2-cyanoacetate and elemental sulfur, was subjected to a *one-pot* Gewald reaction³⁹ in
5 presence of triethylamine as base to afford ethyl 6-acetyl-2-amino-4,5,6,7-tetrahydrothieno[2,3-
6 c]pyridine-3-carboxylate (CRCM5484-2, **34-2**), in good yields. Reacting this latter with 2-furfuryl
7 isothiocyanate in acetonitrile afforded corresponding 7-acetyl-3-(furan-2-ylmethyl)-2-mercapto-5,6,7,8-
8 tetrahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-4(3H)-one (CRCM5484-1, **34-1**). Finally, condensation
9 with 2-chloro-*N*-(2-methylpyridin-3-yl)acetamide (CRCM5484-3, **34-3**), previously obtained by reaction
10 of appropriate substituted aminopyridine with chloroacetyl chloride, afforded the expected THPTP
11 (CRCM5484, **34**) in 90% yields.




13 **Scheme 1. Reagents and Conditions:** i) TEA, EtOH, reflux, 2h, 74%; ii) 2-furfurylisothiocyanate, K₂CO₃, MeCN, reflux, 12h, 10%;
14 iii) chloroacetyl chloride, TEA, DCM, rt, 5h, 82%; iv) K₂CO₃, acetone, reflux, 2h, 90%.

15 **CRCM5484, a BDII selective BET inhibitor**

16 CRCM5484 (Figure 3A) exhibited a BDII selective profile with IC₅₀ values of 130 nM, 20 nM and 71 nM
17 for the BDIIs and 1300 nM, 9500 nM and 4700 nM for the BDIs of BRD4, BDR3 and BRD2, respectively
18 (Figure 3B). The affinity for BRD3-BDII was confirmed by isothermal titration calorimetry (ITC) with a 150
19 nM measured K_D (Figure 3C). The BROMOscan evaluation at 1μM confirmed the BDII selectivity profile
20 (Figure 3D and Table S1) and also an overall BET family selectivity across the 32 evaluated domains.

1 From this set of data, only two additional relatively lower affinity binders could be highlighted i.e.
2 TRIM33 (52%) and CECR2 (54%) as compared to the 36, 37 and 41% inhibition measured for the BDII of
3 BRD3, BRD2, and BRD4, respectively. A potential impact of these additional binders, either beneficial or
4 detrimental, cannot be ruled out when assessing the biological effects of CRCM5484. The co-crystal
5 structure of CRCM5484 in complex with BRD4-BDI and BRD2-BDII was obtained at 1.2 and 1.5 Å
6 resolution, respectively (Figure 3E). As expected from the SAR study, the binding mode of CRCM5484 is
7 different than that of the thienopyrimidine derivative compound in the 4ZW1 structure (Figure S3B). For
8 CRCM5484, the N-acetyl from the tetrahydropyridinyl moiety is anchored in the binding pocket and
9 mimics the acetylated lysine through direct hydrogen bonds with the key asparagine residues N140 and
10 N429 in BDI and BDII, respectively. The acetamido group also stabilizes the water network at the bottom
11 of the pocket. A major conformational change is also observed for CRCM5484 between the two BDs, as
12 shown in Figure 3E and Figure S5. The picoline bound to BRD4-BDI (marked with an orange star Figure
13 3F) and the furan to BRD2-BDII (marked with a blue star Figure 3G) showed poor electronic density and
14 their positioning were driven by geometry and residual electron density during refinement. In the BDII
15 structure, the picoline moiety occupies the ZA channel and establishes vdW interactions with the WPF
16 shelf residues W370, P371 and F372. This position is also stabilized by various interactions with water
17 molecules and other protein residues. In BDI, this moiety is oriented toward the Y139 and N140 amino
18 acids (Figure 3F and Figure S5A and E). The furyl moiety that we originally linked to the BDII selective
19 profile does not establish direct interactions with H433, the main hot spot for the development of BDII
20 selective drugs (Figure 3G and Figure S5C and F). The crystal structure links the BDII selectivity to a
21 combination of direct or water-mediated anchor points that stabilize the compound in the cavity. In
22 both structures, the furyl moiety binds a pocket formed by the WPF shelf and an aspartate residue
23 (D145 for BRD4-BDI and D434 for BRD2-BDII). Altogether, these results confirm the originality of the


- 1 THPTP molecular mode of action in the BD cavity and its potential as a chemical platform for the
- 2 development of potent BD inhibitors.



3
 4 **Figure 3. Characterization of CRCM5484.** (A) Chemical structure. (B) HTRF selectivity assay on the BDI (solid lines) and BDII
 5 (dashed lines) of BRD2 (blue), BRD3 (green) and BRD4 (red). (C) Representative Isothermal Titration Calorimetry (ITC)
 6 experiment with CRCM5484 in the cell and BRD3-BDII in the syringe at 15°C. (D) BROMOscan data (DiscoverX) on the 8 BET
 7 bromodomains families. (E) Superposition of the co-crystal structures of CRCM5484 (orange)/BRD4-BDI (red) (PDB: 7Q3F) and
 8 CRM5484 (cyan) / BRD2-BDII (blue) (PDB: 7Q50). Water molecules are shown as red and blue spheres for the BRD4-BDI and
 9 BRD2-BDII complexes, respectively. F and G) Co-Crystal structures of CRCM5484 with BRD4-BDI (Orange) and BRD2-BDII (Blue)
 10 showing the hydrogen bond interactions as green dotted lines, the canonical asparagine binding residue as magenta, the WPF
 11 shelf as green and the rest of the domain as white balls and sticks. The picoline (orange *) and furan moiety (blue *) showed
 12 poor electronic density and their positioning were driven by geometry and residual electron density during refinement.

13 **CRCM5484 has similar biological effects as other BDII selective compounds in cell-based assays.**

1 The biological effects of CRCM5484 were evaluated and compared to the reference BDII selective
2 compounds RVX-208 and GSK046, and the pan-BET inhibitor OTX-015. First, treatment of the MOLM-14
3 leukemic cell line showed that CRCM5484, similar to RVX-208 and GSK046, exhibited >100x lower
4 impact on cell growth and viability than that of OTX-015, with EC₅₀ values of 8.5 μM, 6.7 μM, 14 μM and
5 0.08 μM, respectively (Figure 4A). Second, the transcriptional impact of these various BD inhibitors on
6 *MYC* gene expression⁴⁰ was compared by quantitative RT-PCR. All tested molecules induced a
7 concentration-dependent decrease in *MYC* gene expression, with the pan-BET inhibitor displaying the
8 highest potency (IC₅₀ = 0.6 μM), and the BDII selective compounds were at least 5x less efficient with IC₅₀
9 values ranging between 3 and 7 μM (CRCM5484) (Figure 4B). These evaluations were replicated in three
10 additional leukemia cell lines, the known bromodomain inhibitor sensitive CCRF-CEM in addition to
11 MOLM-14 and the two resistant cell lines, NB4 and K-562. Overall, CRCM5484 and the two other
12 evaluated BDII selective compounds (RVX-208 and GSK046) displayed little to no cytotoxicity on the 4
13 cell lines with EC₅₀ > 5μM, however MOLM-14 and CCRF-CEM cells appeared slightly more responsive to
14 the drugs (Figure S6A). *MYC* gene expression profile was also similar for these cell lines (Figure S6B). As
15 BDII inhibition was recently shown to modulate the inflammatory stimulation of gene transcription in
16 MHC-I antigen presentation pathway gene components²⁵, we also evaluated the impact of CRCM5484
17 on K-562 leukemic cells stimulated with the proinflammatory cytokine interferon-γ (IFN_γ). As recently
18 reported, the pan-BET inhibitor totally prevented IFN_γ-induced expression of MHC-I while CRCM5484
19 partially inhibited this induction in a concentration-dependent manner, with a similar efficiency to
20 GSK046 and in agreement with Gilan et al³⁰ (Figure 4C). Although it cannot be excluded that the two
21 molecules hold distinct cell penetration capabilities, these results show that CRCM5484 exhibits
22 comparable activity to the compound GSK046 in these cellular assays.




1

2 **Figure 4. CRCM5484 compared to other BDII selective compounds.** (A) Cell viability assay on MOLM-14 leukemic cells
 3 performed after 72h incubation with OTX015 (black), RVX-208 (blue), GSK046 (green) and CRCM5484 (red). (B) Quantification
 4 of MYC expression by RT-qPCR following 2h incubation with OTX015 (black), RVX-208 (blue), GSK046 (green) and CRCM5484
 5 (red). (C) Flow cytometry analysis of MHC-I expression in K-562 cells following stimulation by IFN γ and treatment with DMSO,
 6 OTX015, GSK046 or CRCM5484.

7 **Cell- and context-dependent modulation of drug sensitivity of leukemic cells by CRCM5484**

8 Because of the limited impact of BDII selective compounds as single agents in cell viability/growth
 9 assays, we next questioned the potential to use them in combination with other known compounds
 10 demonstrating anti-leukemic activity. Using our previously described DSRP platform⁴¹, CRCM5484 was
 11 added to 3 distinct leukemia cell lines (MOLM-14, U-937 and CCRF-CEM), as well as to primary patient-
 12 derived xenografted (PDX) acute myeloid leukemia cells at its 40% effective concentration (EC_{40})
 13 (previously established for each individual cell line and PDX-derived cell cultures, see Table S2 for
 14 details), in combination with a panel of 78 drugs. Since the biological impact of selective BDII inhibition
 15 is best evidenced in inflammatory conditions and because drug sensitivity and resistance profiles of
 16 primary leukemia cell cultures have been shown to be differentially modulated in the presence of bone-


1 marrow-derived stromal cell conditioned medium (BMSC-CM)⁴², PDX-derived leukemia cells⁴³ were
2 cultured in both the absence and presence of BMSC-CM. The fold potentiation ratio was then calculated
3 as the ratio of each tested drug EC₅₀ value determined in the presence and absence of CRCM5484
4 (Figure 5A and B). By arbitrarily setting a threshold of potentiation of at least 3-fold, CRCM5484
5 potentiated the cell viability/growth inhibitory activity of 16 drugs: 10 in MOLM-14, 1 in U-937, 3 in the
6 CCRF-CEM cell line and 2 in the PDX-derived leukemia cells in the absence of BMSC-CM. CRCM5484-
7 potentiated drugs varied with the investigated cell cultures, since only 1 out of these 16 drugs, namely
8 birinapant, was potentiated in at least 3 out of the 4 distinct cell cultures, including PDX-derived
9 leukemia cells. Strikingly, adding BMSC-CM to the PDX-derived leukemia cell cultures greatly modified
10 the drug sensitivity and resistance profiles; 7 additional drugs were potentiated >3-fold by CRCM5484
11 (Figure 5B), but none of these drugs were also potentiated in the established leukemia cell lines (Figure
12 5A).



1
2 **Figure 5. CRCM5484 potentiation of pharmacological agents.** (A) Fold of drug cytotoxic effects with or without CRCM5484 at
3 its EC₄₀ (Table S2) in CCRF-CEM (green), U-937 (blue) and MOLM-14 (red) cell lines. (B) Fold of drug cytotoxic effects with or
4 without CRCM5484 with or without addition of bone-marrow-derived stromal cell conditioned medium (BMSC-CM).

5 To confirm these results on selected drug combinations, we developed dose-response experiments
6 with combinatorial effects analyses on PDX-derived leukemia cells. First, we focused on birinapant as
7 the sole drug identified in our screening strategy in 3 out of 4 cell cultures, including PDX-derived
8 leukemia cells cultured in the absence of BMSC-CM. As shown in figure 6A, a dose-dependent but
9 limited potentiation of birinapant cell viability/growth inhibitory activity by CRCM5484 was observed,
10 which was scored as mild additivity to low synergy by Combenefit⁴⁴ matrix analysis (Figure 6B). Out of
11 the 7 drugs in which the anti-leukemic activity was potentiated by CRCM5484 in the presence of BMSC-
12 CM, we focused on vorinostat. Indeed, for four of these drugs (nelarabine, quizartinib, sorafenib and
13 gefitinib), the potentiation by CRCM5484 most likely resulted from increased resistance induced by
14 BMSC-CM, as previously reported⁴². Furthermore, among the remaining three drugs (vorinostat,
15 masitinib and SG1-110), vorinostat displayed the highest potentiation ratio (11-fold) and has been

1 previously described to synergize with pan-BET inhibitors⁴⁵. Dose-response experiments with
 2 combinatorial effect analyses confirmed the absence of vorinostat potentiation by CRCM5484 in the
 3 absence of BMSC-CM (Figure 6C and D) and dose-dependent potentiation in the presence of BMSC-CM
 4 (Figure 6E and F), which was scored as synergistic by Combenefit matrix analysis at a range of vorinostat
 5 concentrations that are clinically achievable⁴⁶.
 6 Altogether, these results support the differential ability of CRCM5484 to potentiate anti-leukemic
 7 drug activity in a cell and culture condition-dependent manner.



8
 9 **Figure 6. Validation of CRCM5484 potentiation of pharmacological agents in conditioned medium in the presence/absence of**
 10 **BMSC-CM.** Dose response curves of birinapant (A) or vorinostat (C and E) cytotoxic effect at a concentration range of
 11 CRCM5484 in the absence (A and C) or the presence (E) of BMSC-CM. Combination effect matrix between birinapant (B) and
 12 vorinostat (D and F) in combination with CRCM5484 analyzed using the BLISS (right) and HSA model (left) in the absence (B and
 13 D) or the presence (F) of BMSC-CM. The synergy/antagonism score and its statistical significance is specified for each
 14 combination. Significant combinations are colored in an antagonism/synergy scale (red to blue) and stars indicate the level of
 15 significance.

1 DISCUSSION AND CONCLUSIONS

2 The development of selective BDI and/or BDII compounds has been strongly motivated by the
3 encouraging results of first-generation pan-BET inhibitors in various clinical trials, which have even led to
4 remissions in some cancer patients⁴⁷⁻⁴⁹. However, the initial enthusiasm was rapidly tempered by the
5 short response durations observed at tolerated doses and the description of resistance mechanisms that
6 may contribute to these limitations. Therefore, developing more selective molecules was quickly
7 proposed as well as evaluating combinations with standard treatments and/or other targeted therapies
8 to reduce side effects while improving efficacy. Designing inhibitors with a selective profile could also
9 lead to a better understanding of the role played by each domain in the BD family, which is currently
10 under debate, and to propose new original chemistry-based molecules that could be developed as drug
11 candidates.

12 In this study, we describe how a differential screen of BDI and BDII domains of the BET family against
13 our in-house “Fr-PPICChem” chemical library led to the identification of a BD binder with a preference for
14 BDII of the BRD3 and BRD2 proteins from the BET family. This compound has an original selectivity
15 profile based on a THPTP core and has not been previously described as a BD binder in the literature. To
16 optimize this screening hit, a series of five compounds was first purchased during SAR by a catalog
17 campaign, and the molecular mode of action of the best compounds was investigated. The acetamido
18 group appeared to be crucial for compound anchoring in the binding pocket while the furyl moiety was
19 responsible for the selectivity profile.

20 In a second series of nineteen compounds, phenyl substitution could adjust the affinity of the
21 compounds to BDs, and we observed that substitutions in the -Para position, such as ethoxy, methoxy,
22 methyl, chloro or fluoro groups decreased compound affinity with an $IC_{50} > 500$ nM on BRD3-BDII. We
23 predicted that the best compounds of the series established additional hydrogen bonds with a hydrogen
24 bond donor or acceptor in the -Ortho position on the phenyl ring. Within this series, compound **16**

1 exhibited the best inhibitory potential with IC₅₀ values measured at 2200 and 72 nM for the BDI and BDII
2 of BRD3 respectively, and a 31-fold selectivity ratio.

3 A last step in the chemical synthesis aimed at improving the compound selectivity and solubility and
4 led to CRCM5484, our lead compound that bears a picoline moiety instead of the initial aryl moiety. This
5 compound is a BDII selective inhibitor that exhibits a similar activity in TR-FRET to GSK046 or RVX-208,
6 the reference BDII selective compounds in the literature. Comparative analysis of the X-Ray structures of
7 CRCM5484 in complex with BRD4-BDI and BRD2-BDII supported the observations from the SAR by
8 catalog and revealed that the compound was rotated by 180° in the cavity. In BRD4-BDI, the picoline
9 moiety established no interaction with the protein, while in BDII, the picoline moiety was stabilized by
10 vdW contacts in the protein WPF shelf region.

11 We then evaluated the activity of our THPTP-based BDII inhibitor in a variety of cell-based assays, in
12 comparison with two reference BDII selective inhibitors: the recently developed GSK046³⁰, and RVX-208,
13 the earliest described BDII compound¹⁴. Consistent with the activities reported for the latter
14 compounds, when used as a single agent, CRCM5484 presented a weak impact on the
15 viability/proliferation of the leukemic MOLM-14 cell line (as well as U-937 and CCRF-CEM) with EC₅₀
16 values >100x compared to those of the pan-BET compound OTX-015, as well as a limited impact on the
17 maintenance of *MYC* gene transcription, a well-documented transcriptional target of BET inhibitors.

18 In parallel, the work of Gilan *et al* demonstrated the contributions of BDII to the transcriptional
19 induction of genes, and BDII notably controlled the cell surface expression of the MHCI complex by
20 inflammatory cytokines and its modulation by the BDII selective compound GSK046. Consistent with
21 these observations, our results demonstrate that CRCM5484 modulates IFN_γ-induced MHCI expression
22 in a manner comparable to that of GSK046 but is less efficient than the pan-BET inhibitor OTX-015.

23 Based on these observations, we then assessed the ability of CRCM5484 to potentiate the anti-
24 leukemic activity of a panel of 78 drugs with marketing authorization and/or in clinical development.

1 Our screening approach highlights several observations. First, when CRCM5484 is added in a single dose
2 corresponding to its previously measured EC₄₀ in each of the cell models evaluated, the antiproliferative
3 activity of the tested drugs was potentiated up to 9-fold, depending on the cell line and including
4 primary AML cells derived from a PDX. Second, the drugs potentiated by CRCM5484 varied greatly
5 depending on the cell line evaluated and the presence of inflammatory conditioned medium from bone
6 marrow stromal cells for the test performed on primary PDX-derived AML cells (up to 17-fold increased
7 antiproliferative activity for the later). Third, the selection of the two drugs with the best combination
8 potential confirmed the potentiation effect of CRCM5484 in dose-response experiments, which were
9 performed on primary PDX-derived AML cells with additivity/low synergy effects in combination with
10 birinapant and synergy effects in combination with vorinostat, in the presence of conditioned medium
11 from bone marrow stromal cells.

12 To our knowledge, these results are the first illustration of a combination effect of BET inhibitors with
13 birinapant, a second mitochondrial-derived activator of caspases (SMAC) mimetic, in which the rationale
14 for the combination effect remains to be unveiled and better documented. It should be noted that
15 although this potentiation was observed for two of the three cell lines as well as the primary AML cells
16 from our PDX model, it was not observed for the latter in the presence of BMSC-CM.

17 The potentiation effect of the histone deacetylase inhibitor vorinostat by pan-BET inhibitors has
18 recently been reported in cutaneous T-cell lymphoma⁵⁰. However, it is important to note that this effect
19 was only observed for primary AML cells from our PDX model and in the presence of BMSC-CM. It
20 therefore seems important to confirm these observations and in particular to extend them to a panel of
21 primary AML samples.

22 Nevertheless, our results provide motivation to further explore the capacities of CRCM5484 and, more
23 broadly, selective BDII inhibitors to potentiate the anti-leukemic activity of FDA-approved drugs or

1 molecules in development; and the inhibitors should be examined in terms of the potentially reduced
2 side effects due to their better selectivity in various preclinical cancer models.

3 EXPERIMENTAL SECTION

4 **Chemistry**

5 For chemical synthesis, all compounds are >95% pure by HPLC analysis. The Table S3 summarizes all
6 compounds information (SMILES, CAS, provider, and provider ID). Quality evaluations as provided by the
7 company (NMR/HPLC) for all purchased compounds are included as supplementary files. Commercially
8 available reagents and solvents were used without further additional purification. Thin layer
9 chromatography (TLC) was performed on precoated aluminum sheets of silica (60 F254 nm, Merck) and
10 visualized using short-wave UV light. Reaction monitoring and purity of synthesized compounds were
11 recorded by using analytical Agilent Infinity high performance liquid chromatography (HPLC) with DAD at
12 254 nm column Agilent Poroshell 120 EC-C18 2.7 μ m (4.6 x 50 mm), mobile phase (A: 0.1% FA H₂O, B:
13 0.1% FA MeCN), flow rate 0.3 mL/min, time/%B 0/10, 4/90, 7/90, 9/10, 10/10. Column chromatography
14 was performed on a Reveleris purification system using Reveleris Flash silica cartridges. Petroleum refers
15 to the fraction with distillation range 40-65 °C. ¹H and ¹³C NMR spectra were recorded by using a Bruker
16 AC 400 spectrometer. Chemical shifts, (δ) are reported in ppm and coupling values (*J*) in hertz.
17 Abbreviations for peaks are, br: broad, s: singlet, d: doublet, t: triplet, q: quadruplet, quint: quintuplet,
18 sext: sextuplet, sept: septuplet and m: multiplet). The spectra recorded are consistent with the
19 proposed structures. Low-resolution mass spectra were obtained with Agilent SQ G6120B mass
20 spectrometer in positive and negative electrospray mode.

21 **Ethyl 6-acetyl-2-amino-4,5,6,7-tetrahydrothieno[2,3-c]pyridine-3-carboxylate (CRCM5484-2, 34-2)⁵¹.**

22 To a suspension of ethyl 2-cyanoacetate (5.66 g, 50 mmol), *N*-acetyl-4-piperidinone (5.71 g, 50 mmol),
23 triethylamine (7 mL, 50 mmol) and sulfur (1.60 g, 50 mmol) in ethanol (100 mL) was heated at 70 °C for
24 2 h. The resulting mixture was concentrated under reduced pressure and the residue was successively

1 diluted with DCM (100 mL), washed with H₂O (2 x 30 mL), brine (30 mL), and dried over Na₂SO₄. The
2 solvent was distilled off under reduced pressure and the crude product was purified by crystallization
3 from Et₂O to afford ethyl 6-acetyl-2-amino-4,5,6,7-tetrahydrothieno[2,3-c]pyridine-3-carboxylate
4 **CRCM5484-2** (10 g, 75%), as a yellow powder. ¹H NMR (400 MHz, CDCl₃) δ 4.71 (brs, 2H), 4.52-4.37 (m,
5 2H), 4.26 (q, *J* = 7.2 Hz, 2H), 3.79-3.65 (m, 2H), 2.86 (brs, 2H), 2.17 (brs, 3H), 1.36 (t, *J* = 7.2 Hz, 3H); LCMS
6 C₁₂H₁₆N₂O₃S Rt = 6.908 min, ESI+ *m/z* = 269.1 (M+H).

7 **7-Acetyl-3-(furan-2-ylmethyl)-2-mercapto-5,6,7,8-tetrahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-**
8 **4(3H)-one (CRCM5484-1, 34-1)**. To a solution of **CRCM5484-2** (5.36 g, 20 mmol) and 2-furfuryl
9 isothiocyanate (2.76 g, 20 mmol) in acetonitrile (50 ml), was added anhydrous potassium carbonate
10 (2.76 g, 20 mmol). The resulting mixture was heated under reflux overnight. After cooling down at room
11 temperature, the precipitate formed was filtered off, and resuspended, under stirring, in 2N HCl
12 aqueous solution (30 mL) for 15 min. The solid was collected by filtration to afford the 7-acetyl-3-(furan-
13 2-ylmethyl)-2-mercapto-5,6,7,8-tetrahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-4(3H)-one
14 **CRCM5484-1** (700 mg, 10%), as a light yellow powder. ¹H NMR (400 MHz, DMSO) δ 13.71 (brs, 1H), 7.54
15 (d, *J* = 1.2 Hz, 1H), 6.37 (dd, *J* = 3.1, 1.2 Hz, 1H), 6.31 (d, *J* = 3.1 Hz, 1H), 5.57 (s, 2H), 4.64 (s, 0.5H), 4.59 (s,
16 1.5H), 3.71-3.67 (m, 2H), 2.94 (t, *J* = 5.4 Hz, 1.5H), 2.81 (t, *J* = 5.4 Hz, 0.5H), 2.10 (s, 2.25H), 2.05 (s,
17 0.75H); LCMS C₁₆H₁₅N₃O₃S₂ Rt = 7.145 min, ESI+ *m/z* = 362.1 (M+H).

18 **2-Chloro-N-(2-methylpyridin-3-yl)acetamide (CRCM5484-3, 34-3)**⁵². At room temperature, to a
19 solution of 3-amino-2-methylpyridine (2.16 g, 20 mmol) in dichloromethane (20 mL), were successively
20 added chloroacetyl chloride (1.6 mL, 20 mmol) and triethylamine (2.79 mL, 20 mmol). The resulting
21 mixture was stirred for 5 h, and concentrated under reduced pressure. The residue was purified by
22 column chromatography, eluent DCM-MeOH (80:20) to afford the 2-chloro-*N*-(2-methylpyridin-3-
23 yl)acetamide **34-3**. (3.02 g, 82%), as a white powder. LCMS C₈H₉ClN₂O Rt = 2.179 min, ESI+ *m/z* = 185.1
24 (M+H).

1 **2-((7-Acetyl-3-(furan-2-ylmethyl)-4-oxo-3,4,5,6,7,8-hexahydropyrido[4',3':4,5]thieno[2,3-**
2 **d]pyrimidin-2-yl)thio)-N-(2-methylpyridin-3-yl)acetamide (CRCM5484, 34).** To a suspension of
3 **CRCM5484-1** (181 mg, 0.5 mmol) and anhydrous potassium carbonate (170 mg, 1.23 mmol) in dry
4 acetone (15 mL) was added **CRCM5484-3** (139 mg, 0.75 mmol). The resulting mixture was refluxed for 1
5 h and concentrated under reduced pressure. The residue was purified by column chromatography,
6 eluent DCM-MeOH (95:5) to afford the 2-((7-acetyl-3-(furan-2-ylmethyl)-4-oxo-3,4,5,6,7,8-
7 hexahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidin-2-yl)thio)-N-(2-methylpyridin-3-yl)acetamide
8 **CRCM5484** (230 mg, 90%), as a white powder. CRCM5484 purity was analyzed by HPLC as shown in
9 Figure S7A. ¹H NMR (400 MHz, CDCl₃) δ 8.71 (brs, 1H), 8.36-8.31 (m, 1H), 8.27 (d, *J* = 5.1 Hz, 1H), 7.35
10 (brs, 1H), 7.22 (dd, *J* = 8.1, 5.1 Hz, 1H), 6.45 (d, *J* = 3.2 Hz, 1H), 6.33 (dd, *J* = 3.2, 1.9 Hz, 1H), 5.35 (s, 2H),
11 4.77 (s, 1.5H), 4.64 (s, 0.5H), 4.11 (s, 2H), 3.88 (t, *J* = 5.6 Hz, 0.5H), 3.74 (t, *J* = 5.6 Hz, 1.5H), 3.16 (t, *J* = 5.6
12 Hz, 1.5H), 3.10 (t, *J* = 5.6 Hz, 0.5H), 2.43 (s, 3H), 2.21 (s, 2.25H), 2.19 (s, 0.75H); ¹³C NMR (10 MHz, CDCl₃)
13 δ 169.53, 169.40, 166.59, 166.53, 161.69, 157.39, 157.21, 157.06, 156.86, 149.48, 149.34, 147.70,
14 145.29, 142.81, 132.14, 131.48, 130.29, 130.22, 129.63, 128.77, 126.97, 121.76, 118.72, 118.57, 110.65,
15 110.51, 110.43, 45.23, 43.51, 41.13, 40.53, 38.68, 36.44, 26.14, 25.26, 22.03, 21.42, 20.90; LCMS
16 C₂₄H₂₃N₅O₄S₂ Rt = 6.298 min, ESI+ *m/z* = 510.1 (M+H) as shown in Figure S7B, >98% pure.

17 **Protein expression and purification**

18 For homogeneous time resolved fluorescence (HTRF) experiments, all 6 BDs of BRD4, BRD3 and BRD2
19 (3 BDI and 3 BDII domains) synthetic genes that include a Tobacco Etch Virus (TEV) cleavage site were
20 purchased from LifeTechnology in a pDONR transport vector before cloning into a pDESTTM15 expression
21 vector for GST affinity purification. Protein production and purification was carried out using similar
22 protocols and buffers used for the His-BRD4-BDI system. Purification was carried on GST affinity resin
23 (Thermo Scientific) and reduced glutathione was used for protein release. GST-BRD4-BDI was further

1 purified by size exclusion chromatography on a Superdex 16/60 Hilo column (GE Healthcare) using 20
2 mM TRIS pH=8.0, 150 mM NaCl Buffer.

3 For isothermal titration calorimetry (ITC), BRD3-BDII was produced and purified using a histidine tag
4 affinity chromatography from a pDESTTM17 expression vector containing a TEV protease cleavage site.
5 After size exclusion chromatography, fractions presenting pure BRD3-BDII were pooled and
6 concentrated to 7 mg/mL in 10 mM HEPES pH=7.5, 150 mM NaCl, 1mM DTT buffer. A final step of buffer
7 exchange was performed using a PD10 column from GE healthcare prior to the experiment to remove
8 the DTT.

9 For X-Ray crystallography, BRD4-BDI was produced and purified using a histidine tag affinity
10 chromatography as described by Filipakopoulos et al⁴⁹. For these experiments, a pNIC28-BSA4
11 expression vector containing BRD4-BDI and a TEV protease cleavage site has been kindly provided by
12 Stefan Knapp laboratory from the SGC at the University of Oxford. After size exclusion chromatography,
13 fractions of pure BRD4-BDI after TEV cleavage of the histidine tag were pooled and concentrated to 25
14 mg/mL.

15 For X-Ray crystallography, BRD2-BDII was produced from a pDESTTM17 expression vector containing a
16 TEV protease cleavage site and purified using a histidine tag affinity chromatography. After size
17 exclusion chromatography, fractions of pure BRD2-BDII after TEV cleavage were pooled and
18 concentrated to 20 mg/mL.

19 **Homogeneous Time-Resolved Fluorescence**

20 HTRF assays were performed in white 1536 Well Small VolumeTM HiBase Polystyrene Microplates
21 (Greiner) with a total working volume of 4 μ L using the conditions described in Table S4. 5 nL of
22 compounds were dispensed (2 x 2.5 nL fixed dispensing using an Echo550 (Labcyte)) from a
23 concentration stock of 1 mM (100% DMSO) for the Fr-PPIchem primary screening after addition of 15 nL
24 DMSO per well to reach the final 1.25 μ M concentration used in the assay (0.5% final DMSO). Dose

1 response evaluations of commercial and synthesized compounds were performed similarly from 10 mM
2 stock solutions (100% DMSO). For the screening, the inhibition percentage was calculated and compared
3 to the reference compound, JQ1. The IC₅₀ measurements were carried out in triplicates. All HTRF
4 reagents were purchased from CisBio Bioassays and reconstituted according to the supplier protocols.
5 HTRF signals were measured, after a final incubation (6h at room temperature), using a PHERAstar FS
6 (BMG Labtech) with an excitation filter at 337 nm and fluorescence wavelength measurement at 620
7 and 665 nm, using an integration delay of 60 μs and an integration time of 500 μs. Results were analyzed
8 with a two-wavelengths signal ratio: [intensity (665 nm)/intensity (620 nm)]*10⁴. Percentage of
9 inhibition was calculated using the following equation: % inhibition = [(compound signal) - (min signal)] /
10 [(max signal) - (min signal)] * 100, where 'max signal' is the signal ratio with the compound vehicle alone
11 (DMSO) and 'min signal' the signal ratio without peptide. For IC₅₀ measurements, values were
12 normalized and fitted with Prism (GraphPad) using the following equation: $Y = 100 / (1 + ((X / IC_{50})^{Hill$
13 $slope}))$.

14 **BROMOscan**

15 BROMOscan bromodomain profiling was provided by Eurofin DiscoverX Corp. Determination of the K_D
16 between test compounds and DNA tagged bromodomains was achieved through binding competition
17 against a proprietary reference immobilized ligand. CRCM5484 was used at 1μM.

18 **Isothermal Titration Calorimetry**

19 ITC was used to evaluate the thermodynamics parameters of the binding between BRD4-BDI, BRD2-
20 BDI and BRD3-BD2 and CRCM5484, using ITC conditions previously described by Filippakopoulos et al⁵³.
21 Purified proteins were extensively dialyzed in the ITC buffer containing 10mM Hepes pH=7.5 and 150
22 mM NaCl. Compound was diluted directly in the last protein dialysate prior to experiments. Titrations
23 were carried out on a MicroCal ITC200 microcalorimeter (GE Healthcare, Piscataway, NJ). Each
24 experiment was designed using a titrant concentration (protein in the syringe) set 10 to 15 times the

1 analyte concentration (compound in the cell generally between 10 and 35 μM) and using 13 injections at
2 15 $^{\circ}\text{C}$. A first small injection (generally 0.2 μL) was included in the titration protocol to remove air
3 bubbles trapped in the syringe prior titration and/or take into account syringe predilution in the cell
4 during equilibration. Raw data were scaled after setting the zero to the titration saturation heat value.
5 Integrated raw ITC data were fitted to a one site non-linear least squares fit model using the MicroCal
6 Origin plugin as implemented in Origin 7 (Origin Lab). Finally, ΔG and $\text{T}\Delta\text{S}$ values were calculated from
7 the fitted ΔH and K_A values using the equations $\Delta\text{G} = -\text{R}\cdot\text{T}\cdot\ln\text{K}_\text{A}$ and $\Delta\text{G} = \Delta\text{H} - \text{T}\Delta\text{S}$.

8 **X-Ray crystallography**

9 BRD4-BDI/CRCM5484 and BRD2-BDII/CRCM5484 co-crystallizations were performed at respectively,
10 4 $^{\circ}\text{C}$ (277 K) and 12 $^{\circ}\text{C}$ (285 K) using the hanging drop vapor diffusion method. For the BRD4-
11 BDI/CRCM5484, a solution of 25 mg/mL of BRD4-BDI with 1mM of compound was mixed at a 1:1 ratio
12 with the precipitant solution (300 mM NaNO_3 , 16% PEG3350, 5% Ethylene glycol) and crystals grew to
13 diffracting quality within 3-5 days.

14 For the BRD2-BDII/CRCM5484 complex, 20 mg/mL of BRD2-BDII and 2.5 mM of compound CRCM5484
15 preparation was mixed at a 1:1 ratio with the precipitant solution (25% PEG3350, 100 mM potassium
16 thiocyanate) and crystals grew to diffracting quality within 10 days. Crystals grew to diffracting quality
17 within 1 month.

18 Crystals were cryo-protected using the precipitant solution supplemented with 10% ethylene glycol
19 and were flash frozen in liquid nitrogen. Data were collected at the ESRF beamlines ID30A-1 and at the
20 SOLEIL beamline Proxima1. Indexing, integration, and scaling were performed using XDS. Initial phases
21 were calculated by molecular replacement with Phaser MR (CCP4 suite) using a model of the first
22 domain of BRD4 (extracted from the Protein Data Bank accession code: 2OSS); for BRD2-BDII, using a
23 model extracted from the Protein Data Bank accession code: 5XHK. Initial models for the protein and the
24 ligands were built in COOT. The cycles of refinement were carried out with phenix.refine (Phenix v1.13).

1 Data collection and refinement statistics can be found in Table S5. The models and structures factors
2 have been deposited with Protein Data Bank accession code for BRD4-BDI in complex with compound
3 CRCM5484: 7Q3F, and BRD2-BDII in complex with the compound CRCM5484: 7Q5O.

4 **Cell culture**

5 The human leukemia cell lines U-937, CCRF-CEM and K-562 were cultured in RPMI 1640
6 supplemented with 10% FBS. MOLM-14 were cultured in MEMalpha supplemented with 10% FBS. NB4
7 were cultured in Dulbecco's Modified Eagle Medium supplemented with 10% FBS. All cell lines were
8 incubated at 37°C with 5% CO₂, were diluted every 2 to 3 days and maintained in an exponential growth
9 phase. These cell lines have been routinely tested for mycoplasma contamination.

10 **Cell viability assay**

11 MOLM-14, NB4, CCRF-CEM and K562 were seeded at 4000, 5000, 8000 and 6000 cells/well in 96-well
12 plates, respectively. After 24h, cells were treated with a range of concentrations of CRCM5484, OTX015,
13 RVX208 or GSK046 and after 72h drug incubation, metabolic activity was detected by addition of Alamar
14 blue and spectrophotometric analysis using a PHERAstar plate reader (BMG labtech). Cell viability was
15 determined and expressed as a percentage of untreated control cells. The determination of IC₅₀ values
16 was performed by using the following equation: $Y=100/(1+((X/IC_{50})^{Hillslope}))$.

17 **MYC gene expression analysis by RT-qPCR**

18 MOLM-14 cells were treated with a range of concentrations of CRCM5484, OTX-015, RVX-208 or
19 GSK046 for 2h. NB4, CCRF-CEM and K-562 cells were treated with 1 μM of OTX-015, 5 μM of CRCM5484,
20 5 μM of GSK046 or 5 μM of RVX-208 for 2h. RNA extraction was performed using RNeasy kit (Qiagen).
21 OneScript® Plus cDNA Synthesis Kit (abm) was used to prepare cDNA from purified RNA. Real-time qPCR
22 was performed using SsoAdvanced Universal SYBR® Green Supermix (Bio-Rad) and a CFX384™ Real-
23 Time System device (Bio-Rad). Analysis performed on duplicate PCR data per biological replicate. Four
24 biological replicates were used for DMSO, CRCM5484, GSK046, RVX208 and OTX015 treated cells. *MYC*

1 gene expression level was determined using the $\Delta\Delta C_t$ method, normalized to *GAPDH* control gene, using
2 pre-designed KiCqStart™ primers (H_MYC_1, H_GAPDH_1; Merck).

3 Inhibition of MHC expression

4 The MHC-I expression was carried out by fluorescence-activated cell sorting analysis using antibodies
5 against HLA-A/B/C (HLA-ABC Monoclonal Antibody (W6/32), PE-Cyanine5, eBioscience) in K-562 cells
6 after stimulation with IFN- γ (10 ng/ml, PHC4031 life technologies) and treatment with DMSO, OTX-015,
7 CRCM5484, or GSK046 for 48 hours. Following incubation and washing, samples were analyzed on a LSR
8 II flow cytometer (Becton Dickinson) using flowjo V-10.8.1 software.

9 **Drug sensitivity and resistance assays**

10 The drug screening library included 78 substances consisting of conventional chemotherapeutics and
11 a broad range of targeted oncology compounds. Each drug was plated on 96-well plates at 4
12 concentrations covering a 1000-fold concentration range. 10-20000 cells were seeded per well in 96-
13 well plates and incubated in the presence of compounds (4 concentrations generated after a 1 log serial
14 dilution from the starting concentration defined in Table S2 for each drug/cell line) in presence or
15 absence of CRCM5484 in a humidified environment at 37°C and 5% CO₂. After 72h (cell lines) - 144h
16 (PDX) of treatment, cell viability was measured using the CellTiter-Glo Luminescent Cell Viability Assay as
17 described by the manufacturer (Promega Corporation). The Luminescence was measured using a Centro
18 luminometer LB960 (Berthold). Effective half-maximal concentration values (EC₅₀) were deduced from
19 dose-response curves obtained using GraphPad Prism 6 software (GraphPad Software, Inc.). Primary
20 AML cells were collected after two passages in mice from a previously described Patient-Derived
21 Xenograft mouse model⁴³ and were cultured with RPMI 20% FBS, 50% bone-marrow-derived stromal cell
22 conditioned medium (HS-5 Cell line) or RPMI 20% FBS, as described³⁹.

23 **Synergy assays**

1 Median effect analysis was performed through Combenefit software (Bioinformatics Advance Access
2 published May 28, 2016) based on Chou and Talalay method using growth inhibition obtained from
3 CellTiter-Glo luminescent assay. The combination effect was compared to the cytotoxicity of each drug
4 alone. We performed combination analyses using Bliss and HSA methods. The drug interactions were
5 specified using Matrix Synergy Plot. The synergy assays were carried out in triplicates

6 **ASSOCIATED CONTENT**

7 **Supporting Information**

8 See figures S1 to S7 and Tables S1 to S5:

9 Figure 1: HTRF evaluation of the FR-PPICChem screening hits, Figure 2: chemical structures, Figure 3:
10 binding mode comparison between CRCM5484 and 4ZW1, Figure 4: selectivity evaluation of compound
11 **12** and compound **16**, Table 1: BROMOScan evaluation of CRCM5484, Figure 5: binding mode study of
12 CRCM5484, Table 2: maximal concentration used for the 78 evaluated compounds in the drug sensitivity
13 and resistance profiling; Figure 6: CRCM5484 HPLC analysis; Table 3: Compounds SMILES, Provider,
14 Provider ID and CAS number; Table 4: HTRF experimental conditions, Table 5: X-ray data collection and
15 refinement statistics. Quality evaluations as provided by the company (NMR/HPLC) are included as
16 supplementary files (file Carrasco-et-al-Providers.zip) for all purchased compounds. CSV file listing
17 compounds formula strings with their associated biochemical and biological data (Carrasco-et-al-
18 Data.csv). This material is available free of charge via the Internet at <http://pubs.acs.org>.

19 **AUTHOR INFORMATION**

20 **Corresponding Authors**

21 *Email: stephane.betzi@inserm.fr. Phone: +33 (0)4 86 97 73 33. ORCID Number: 0000-0001-5935-

22 5058

23 *Email: yves.collette@inserm.fr. Phone: +33 (0)4 86 97 73 31. ORCID Number: 0000-0001-5359-7099

1 *Email: xavier.morelli@inserm.fr. Phone: +33 (0)4 86 97 73 31. ORCID Number: 0000-0001-8101-7901

2 **Author Contributions**

3 K.C. and C.D. carried out the small molecules screen; K.C. and S.B. performed the biochemical and
4 biophysical evaluation; Chemical synthesis was performed by S.C. and J.C.; P.R. and L.H. performed the
5 chemoinformatics and molecular modeling analysis; C.M., K.C., E.P. and A.R. carried out the cellular
6 evaluations. The manuscript was written through contributions of all authors. S.B., Y.C. and X.M.
7 supervised and conceived the project. All authors have given approval to the final version of the
8 manuscript.

9 **PDB ID CODES**

10 PDB codes for BRD4-BDI and BRD2-BDII with bound CRCM5484 are 7Q3F and 7Q5O. Authors will
11 release the atomic coordinates upon article publication.

12 **ACKNOWLEDGMENTS**

13 This study was partly supported by research funding from the "Canceropôle PACA", "Institut National
14 du Cancer" and "Région Provence-Alpes-Côte d'Azur" ("Prematuration Grant" 2017 attributed to X.M. &
15 "Emergence Grant" 2017 attributed to E.P) and from the "Fondation ARC" (PJA 20171206125 grant
16 attributed to X.M.), K.C. was supported by a fellowship from the "Fondation ARC". We would like to
17 thank the FRISBI ANR-10-INSB-05-01 grant for access time on the structural-biology platform of the
18 AFMB laboratory. We acknowledge the European Synchrotron Radiation Facility and the SOLEIL
19 synchrotron for provision of synchrotron radiation facilities and the staff at beamline ID30A-1, Proxima1
20 and Proxima2. We thank the HiTS, TrGET and flow cytometry platforms of the CRCM and their staff. We
21 thank Dr Etienne Rebuffet from iSCB team for fruitful discussions about our structural data.

22

1 **ABBREVIATIONS**

2 BET, Bromodomain and Extra-Terminal; BD, Bromodomain; PPI, Protein-Protein Interactions; ITC,
3 Isothermal Titration Calorimetry; PAINS, Pan-Assay Interference Compounds; HTS, High-Throughput
4 Screen; THPTP, tetrahydropyridothienopyrimidinone; SAR, Structure Activity Relationship; DSRP, Drug
5 Sensitivity Response Profile; HTRF, Homogeneous Time-Resolved Fluorescence; IFN_γ, Pro-Inflammatory
6 Cytokine Interferon-_γ; BMSC-CM, Bone-Marrow-Derived Stromal Cell Conditioned Medium; HPLC, High
7 Performance Liquid Chromatography.

8

1 REFERENCES

- 2 (1) Fujisawa, T.; Filippakopoulos, P. Functions of Bromodomain-Containing Proteins and Their Roles in
3 Homeostasis and Cancer. *Nat Rev Mol Cell Biol* **2017**, *18* (4), 246–262.
- 4 (2) Tough, D. F.; Tak, P. P.; Tarakhovsky, A.; Prinjha, R. K. Epigenetic Drug Discovery: Breaking through
5 the Immune Barrier. *Nat Rev Drug Discov* **2016**, *15* (12), 835–853.
- 6 (3) Tough, D. F.; Prinjha, R. K. Immune Disease-Associated Variants in Gene Enhancers Point to BET
7 Epigenetic Mechanisms for Therapeutic Intervention. *Epigenomics* **2017**, *9* (4), 573–584.
- 8 (4) Bandukwala, H. S.; Gagnon, J.; Togher, S.; Greenbaum, J. A.; Lamperti, E. D.; Parr, N. J.;
9 Molesworth, A. M. H.; Smithers, N.; Lee, K.; Witherington, J.; Tough, D. F.; Prinjha, R. K.; Peters, B.;
10 Rao, A. Selective Inhibition of CD4+ T-Cell Cytokine Production and Autoimmunity by BET Protein
11 and c-Myc Inhibitors. *PNAS* **2012**, *109* (36), 14532–14537.
- 12 (5) Schilderink, R.; Bell, M.; Reginato, E.; Patten, C.; Rioja, -->Inmaculada; Hilbers, F. W.; Kabala, P. A.;
13 Reedquist, K. A.; Tough, D. F.; Tak, P. P.; Prinjha, R. K.; de Jonge, W. J. BET Bromodomain Inhibition
14 Reduces Maturation and Enhances Tolerogenic Properties of Human and Mouse Dendritic Cells.
15 *Molecular Immunology* **2016**, *79*, 66–76.
- 16 (6) Belkina, A. C.; Nikolajczyk, B. S.; Denis, G. V. BET Protein Function Is Required for Inflammation:
17 Brd2 Genetic Disruption and BET Inhibitor JQ1 Impair Mouse Macrophage Inflammatory
18 Responses. *The Journal of Immunology* **2013**, *190* (7), 3670–3678.
- 19 (7) Nicodeme, E.; Jeffrey, K. L.; Schaefer, U.; Beinke, S.; Dewell, S.; Chung, C.; Chandwani, R.; Marazzi,
20 I.; Wilson, P.; Coste, H.; White, J.; Kirilovsky, J.; Rice, C. M.; Lora, J. M.; Prinjha, R. K.; Lee, K.;
21 Tarakhovsky, A. Suppression of Inflammation by a Synthetic Histone Mimic. *Nature* **2010**, *468*
22 (7327), 1119–1123.
- 23 (8) Zhang, Q.; Qian, J.; Zhu, Y. Targeting Bromodomain-Containing Protein 4 (BRD4) Benefits
24 Rheumatoid Arthritis. *Immunology Letters* **2015**, *166* (2), 103–108.
- 25 (9) Prinjha, R. K.; Witherington, J.; Lee, K. Place Your BETs: The Therapeutic Potential of
26 Bromodomains. *Trends in Pharmacological Sciences* **2012**, *33* (3), 146–153.
- 27 (10) Filippakopoulos, P.; Qi, J.; Picaud, S.; Shen, Y.; Smith, W. B.; Fedorov, O.; Morse, E. M.; Keates, T.;
28 Hickman, T. T.; Felletar, I.; Philpott, M.; Munro, S.; McKeown, M. R.; Wang, Y.; Christie, A. L.; West,
29 N.; Cameron, M. J.; Schwartz, B.; Heightman, T. D.; La Thangue, N.; French, C. A.; Wiest, O.; Kung,
30 A. L.; Knapp, S.; Bradner, J. E. Selective Inhibition of BET Bromodomains. *Nature* **2010**, *468* (7327),
31 1067–1073.
- 32 (11) Chaidos, A.; Caputo, V.; Gouvedenou, K.; Liu, B.; Marigo, I.; Chaudhry, M. S.; Rotolo, A.; Tough, D.
33 F.; Smithers, N. N.; Bassil, A. K.; Chapman, T. D.; Harker, N. R.; Barbash, O.; Tummino, P.; Al-Mahdi,
34 N.; Haynes, A. C.; Cutler, L.; Le, B.; Rahemtulla, A.; Roberts, I.; Kleijnen, M.; Witherington, J. J.; Parr,
35 N. J.; Prinjha, R. K.; Karadimitris, A. Potent Antimyeloma Activity of the Novel Bromodomain
36 Inhibitors I-BET151 and I-BET762. *Blood* **2014**, *123* (5), 697–705.
- 37 (12) da Motta, L. L.; Ledaki, I.; Purshouse, K.; Haider, S.; De Bastiani, M. A.; Baban, D.; Morotti, M.;
38 Steers, G.; Wigfield, S.; Bridges, E.; Li, J.-L.; Knapp, S.; Ebner, D.; Klamt, F.; Harris, A. L.; McIntyre, A.
39 The BET Inhibitor JQ1 Selectively Impairs Tumour Response to Hypoxia and Downregulates CA9
40 and Angiogenesis in Triple Negative Breast Cancer. *Oncogene* **2017**, *36* (1), 122–132.
- 41 (13) Rhyasen, G. W.; Hattersley, M. M.; Yao, Y.; Dulak, A.; Wang, W.; Petteruti, P.; Dale, I. L.; Boiko, S.;
42 Cheung, T.; Zhang, J.; Wen, S.; Castriotta, L.; Lawson, D.; Collins, M.; Bao, L.; Ahdesmaki, M. J.;
43 Walker, G.; O'Connor, G.; Yeh, T. C.; Rabow, A. A.; Dry, J. R.; Reimer, C.; Lyne, P.; Mills, G. B.;
44 Fawell, S. E.; Waring, M. J.; Zinda, M.; Clark, E.; Chen, H. AZD5153: A Novel Bivalent BET
45 Bromodomain Inhibitor Highly Active against Hematologic Malignancies. *Mol Cancer Ther* **2016**, *15*
46 (11), 2563–2574.
- 47 (14) Picaud, S.; Wells, C.; Felletar, I.; Brotherton, D.; Martin, S.; Savitsky, P.; Diez-Dacal, B.; Philpott, M.;
48 Bountra, C.; Lingard, H.; Fedorov, O.; Müller, S.; Brennan, P. E.; Knapp, S.; Filippakopoulos, P. RVX-

- 1 208, an Inhibitor of BET Transcriptional Regulators with Selectivity for the Second Bromodomain.
2 *PNAS* **2013**, *110* (49), 19754–19759.
- 3 (15) Filippakopoulos, P.; Picaud, S.; Fedorov, O.; Keller, M.; Wrobel, M.; Morgenstern, O.; Bracher, F.;
4 Knapp, S. Benzodiazepines and Benzotriazepines as Protein Interaction Inhibitors Targeting
5 Bromodomains of the BET Family. *Bioorg Med Chem* **2012**, *20* (6), 1878–1886.
- 6 (16) Noel, J. K.; Iwata, K.; Ooike, S.; Sugahara, K.; Nakamura, H.; Daibata, M. Abstract C244:
7 Development of the BET Bromodomain Inhibitor OTX015. *Mol Cancer Ther* **2013**, *12* (11
8 Supplement), C244–C244.
- 9 (17) Seal, J.; Lamotte, Y.; Donche, F.; Bouillot, A.; Mirguet, O.; Gellibert, F.; Nicodeme, E.; Krysa, G.;
10 Kirilovsky, J.; Beinke, S.; McCleary, S.; Rioja, I.; Bamborough, P.; Chung, C.-W.; Gordon, L.; Lewis, T.;
11 Walker, A. L.; Cutler, L.; Lugo, D.; Wilson, D. M.; Witherington, J.; Lee, K.; Prinjha, R. K.
12 Identification of a Novel Series of BET Family Bromodomain Inhibitors: Binding Mode and Profile of
13 I-BET151 (GSK1210151A). *Bioorg. Med. Chem. Lett.* **2012**, *22* (8), 2968–2972..
- 14 (18) Liu, Z.; Wang, P.; Chen, H.; Wold, E. A.; Tian, B.; Brasier, A. R.; Zhou, J. Drug Discovery Targeting
15 Bromodomain-Containing Protein 4. *J. Med. Chem.* **2017**, *60* (11), 4533–4558.
- 16 (19) Doroshow, D. B.; Eder, J. P.; LoRusso, P. M. BET Inhibitors: A Novel Epigenetic Approach. *Ann Oncol*
17 **2017**, *28* (8), 1776–1787.
- 18 (20) Alqahtani, A.; Choucair, K.; Ashraf, M.; Hammouda, D. M.; Alloghbi, A.; Khan, T.; Senzer, N.;
19 Nemunaitis, J. Bromodomain and Extra-Terminal Motif Inhibitors: A Review of Preclinical and
20 Clinical Advances in Cancer Therapy. *Future Science OA* **2019**, *5*(3) FSO372.
- 21 (21) Andrieu, G.; Belkina, A. C.; Denis, G. V. Clinical Trials for BET Inhibitors Run Ahead of the Science.
22 *Drug Discovery Today: Technologies* **2016**, *19*, 45–50.
- 23 (22) Postel-Vinay, S.; Herbschleb, K.; Massard, C.; Woodcock, V.; Soria, J.-C.; Walter, A. O.; Ewerton, F.;
24 Poelman, M.; Benson, N.; Ocker, M.; Wilkinson, G.; Middleton, M. First-in-Human Phase I Study of
25 the Bromodomain and Extraterminal Motif Inhibitor BAY 1238097: Emerging
26 Pharmacokinetic/Pharmacodynamic Relationship and Early Termination Due to Unexpected
27 Toxicity. *European Journal of Cancer* **2019**, *109*, 103–110.
- 28 (23) Odenike, O.; Wolff, J. E.; Borthakur, G.; Aldoss, I. T.; Rizzieri, D.; Prebet, T.; Hu, B.; Dinh, M.; Chen,
29 X.; Modi, D.; Freise, K. J.; Jonas, B. A. Results from the First-in-Human Study of Mivebresib (ABBV-
30 075), a Pan-Inhibitor of Bromodomain and Extra Terminal Proteins, in Patients with
31 Relapsed/Refractory Acute Myeloid Leukemia. *JCO* **2019**, *37* (15_suppl), 7030–7030.
- 32 (24) Rianjongdee, F.; Atkinson, S. J.; Chung, C.-W.; Grandi, P.; Gray, J. R. J.; Kaushansky, L. J.; Medeiros,
33 P.; Messenger, C.; Phillipou, A.; Preston, A.; Prinjha, R. K.; Rioja, I.; Satz, A. L.; Taylor, S.; Wall, I. D.;
34 Watson, R. J.; Yao, G.; Demont, E. H. Discovery of a Highly Selective BET BD2 Inhibitor from a DNA-
35 Encoded Library Technology Screening Hit. *J Med Chem* **2021**, *64* (15), 10806–10833.
- 36 (25) Seal, J. T.; Atkinson, S. J.; Bamborough, P.; Bassil, A.; Chung, C.-W.; Foley, J.; Gordon, L.; Grandi, P.;
37 Gray, J. R. J.; Harrison, L. A.; Kruger, R. G.; Matteo, J. J.; McCabe, M. T.; Messenger, C.; Mitchell, D.;
38 Phillipou, A.; Preston, A.; Prinjha, R. K.; Rianjongdee, F.; Rioja, I.; Taylor, S.; Wall, I. D.; Watson, R. J.;
39 Woolven, J. M.; Wyce, A.; Zhang, X.-P.; Demont, E. H. Fragment-Based Scaffold Hopping:
40 Identification of Potent, Selective, and Highly Soluble Bromo and Extra Terminal Domain (BET)
41 Second Bromodomain (BD2) Inhibitors. *J Med Chem* **2021**, *64* (15), 10772–10805.
- 42 (26) Lucas, S. C. C.; Atkinson, S. J.; Chung, C.; Davis, R.; Gordon, L.; Grandi, P.; Gray, J. J. R.; Grimes, T.;
43 Phillipou, A.; Preston, A. G.; Prinjha, R. K.; Rioja, I.; Taylor, S.; Tomkinson, N. C. O.; Wall, I.; Watson,
44 R. J.; Woolven, J.; Demont, E. H. Optimization of a Series of 2,3-Dihydrobenzofurans as Highly
45 Potent, Second Bromodomain (BD2)-Selective, Bromo and Extra-Terminal Domain (BET) Inhibitors.
46 *J. Med. Chem.* **2021**, *64* (15), 10711–10741.
- 47 (27) Harrison, L. A.; Atkinson, S. J.; Bassil, A.; Chung, C.-W.; Grandi, P.; Gray, J. R. J.; Levernier, E.; Lewis,
48 A.; Lugo, D.; Messenger, C.; Michon, A.-M.; Mitchell, D. J.; Preston, A.; Prinjha, R. K.; Rioja, I.; Seal,

- 1 J. T.; Taylor, S.; Wall, I. D.; Watson, R. J.; Woolven, J. M.; Demont, E. H. Identification of a Series of
2 N-Methylpyridine-2-Carboxamides as Potent and Selective Inhibitors of the Second Bromodomain
3 (BD2) of the Bromo and Extra Terminal Domain (BET) Proteins. *J Med Chem* **2021**, *64* (15), 10742–
4 10771.
- 5 (28) Gacias, M.; Gerona-Navarro, G.; Plotnikov, A. N.; Zhang, G.; Zeng, L.; Kaur, J.; Moy, G.; Rusinova, E.;
6 Rodriguez, Y.; Matikainen, B.; Vincek, A.; Joshua, J.; Casaccia, P.; Zhou, M.-M. Selective Chemical
7 Modulation of Gene Transcription Favors Oligodendrocyte Lineage Progression. *Chem Biol* **2014**,
8 *21* (7), 841–854.
- 9 (29) Cheung, K.; Lu, G.; Sharma, R.; Vincek, A.; Zhang, R.; Plotnikov, A. N.; Zhang, F.; Zhang, Q.; Ju, Y.;
10 Hu, Y.; Zhao, L.; Han, X.; Meslamani, J.; Xu, F.; Jaganathan, A.; Shen, T.; Zhu, H.; Rusinova, E.; Zeng,
11 L.; Zhou, J.; Yang, J.; Peng, L.; Ohlmeyer, M.; Walsh, M. J.; Zhang, D. Y.; Xiong, H.; Zhou, M.-M. BET
12 N-Terminal Bromodomain Inhibition Selectively Blocks Th17 Cell Differentiation and Ameliorates
13 Colitis in Mice. *Proc Natl Acad Sci U S A* **2017**, *114* (11), 2952–2957.
- 14 (30) Gilan, O.; Rioja, I.; Knezevic, K.; Bell, M. J.; Yeung, M. M.; Harker, N. R.; Lam, E. Y. N.; Chung, C.;
15 Bamborough, P.; Petretich, M.; Urh, M.; Atkinson, S. J.; Bassil, A. K.; Roberts, E. J.; Vassiliadis, D.;
16 Burr, M. L.; Preston, A. G. S.; Wellaway, C.; Werner, T.; Gray, J. R.; Michon, A.-M.; Gobbetti, T.;
17 Kumar, V.; Soden, P. E.; Haynes, A.; Vappiani, J.; Tough, D. F.; Taylor, S.; Dawson, S.-J.; Bantscheff,
18 M.; Lindon, M.; Drewes, G.; Demont, E. H.; Daniels, D. L.; Grandi, P.; Prinjha, R. K.; Dawson, M. A.
19 Selective Targeting of BD1 and BD2 of the BET Proteins in Cancer and Immunoinflammation.
20 *Science* **2020**, *368* (6489), 387–394.
- 21 (31) McLure, K. G.; Gesner, E. M.; Tsujikawa, L.; Kharenko, O. A.; Attwell, S.; Campeau, E.; Wasiaik, S.;
22 Stein, A.; White, A.; Fontano, E.; Suto, R. K.; Wong, N. C. W.; Wagner, G. S.; Hansen, H. C.; Young, P.
23 R. RVX-208, an Inducer of ApoA-I in Humans, Is a BET Bromodomain Antagonist. *PLoS One* **2013**, *8*
24 (12), e83190.
- 25 (32) Law, R. P.; Atkinson, S. J.; Bamborough, P.; Chung, C.; Demont, E. H.; Gordon, L. J.; Lindon, M.;
26 Prinjha, R. K.; Watson, A. J. B.; Hirst, D. J. Discovery of Tetrahydroquinoxalines as Bromodomain
27 and Extra-Terminal Domain (BET) Inhibitors with Selectivity for the Second Bromodomain. *Journal*
28 *of Medicinal Chemistry* **2018**, *61*(10):4317-4334.
- 29 (33) Faivre, E. J.; McDaniel, K. F.; Albert, D. H.; Mantena, S. R.; Plotnik, J. P.; Wilcox, D.; Zhang, L.; Bui,
30 M. H.; Sheppard, G. S.; Wang, L.; Sehgal, V.; Lin, X.; Huang, X.; Lu, X.; Uziel, T.; Hessler, P.; Lam, L.
31 T.; Bellin, R. J.; Mehta, G.; Fidanze, S.; Pratt, J. K.; Liu, D.; Hasvold, L. A.; Sun, C.; Panchal, S. C.;
32 Nicolette, J. J.; Fossey, S. L.; Park, C. H.; Longenecker, K.; Bigelow, L.; Torrent, M.; Rosenberg, S. H.;
33 Kati, W. M.; Shen, Y. Selective Inhibition of the BD2 Bromodomain of BET Proteins in Prostate
34 Cancer. *Nature* **2020**, *578* (7794), 306–310.
- 35 (34) Sheppard, G. S.; Wang, L.; Fidanze, S. D.; Hasvold, L. A.; Liu, D.; Pratt, J. K.; Park, C. H.; Longenecker,
36 K.; Qiu, W.; Torrent, M.; Kovar, P. J.; Bui, M.; Faivre, E.; Huang, X.; Lin, X.; Wilcox, D.; Zhang, L.;
37 Shen, Y.; Albert, D. H.; Magoc, T. J.; Rajaraman, G.; Kati, W. M.; McDaniel, K. F. Discovery of N-
38 Ethyl-4-[2-(4-Fluoro-2,6-Dimethyl-Phenoxy)-5-(1-Hydroxy-1-Methyl-Ethyl)Phenyl]-6-Methyl-7-Oxo-
39 1H-Pyrrolo[2,3-c]Pyridine-2-Carboxamide (ABBV-744), a BET Bromodomain Inhibitor with
40 Selectivity for the Second Bromodomain. *J. Med. Chem.* **2020**, *63* (10), 5585–5623.
- 41 (35) Preston, A.; Atkinson, S.; Bamborough, P.; Chung, C.-W.; Craggs, P. D.; Gordon, L.; Grandi, P.; Gray,
42 J. R. J.; Jones, E. J.; Lindon, M.; Michon, A.-M.; Mitchell, D. J.; Prinjha, R. K.; Rianjongdee, F.; Rioja,
43 I.; Seal, J.; Taylor, S.; Wall, I.; Watson, R. J.; Woolven, J.; Demont, E. H. Design and Synthesis of a
44 Highly Selective and In Vivo-Capable Inhibitor of the Second Bromodomain of the Bromodomain
45 and Extra Terminal Domain Family of Proteins. *J Med Chem* **2020**, *63* (17), 9070–9092.
- 46 (36) Bosc, N.; Muller, C.; Hoffer, L.; Lagorce, D.; Bourg, S.; Derviaux, C.; Gourdel, M.-E.; Rain, J.-C.;
47 Miller, T. W.; Villoutreix, B. O.; Miteva, M. A.; Bonnet, P.; Morelli, X.; Sperandio, O.; Roche, P. Fr-

- 1 PPIChem: An Academic Compound Library Dedicated to Protein–Protein Interactions. *ACS Chem.*
2 *Biol.* **2020**, 15(6):1566–1574.
- 3 (37) Degorce, F.; Card, A.; Soh, S.; Trinquet, E.; Knapik, G. P.; Xie, B. HTRF: A Technology Tailored for
4 Drug Discovery - a Review of Theoretical Aspects and Recent Applications. *Curr Chem Genomics*
5 **2009**, 3, 22–32.
- 6 (38) Chen, Z.; Zhang, H.; Liu, S.; Xie, Y.; Jiang, H.; Lu, W.; Xu, H.; Yue, L.; Zhang, Y.; Ding, H.; Zheng, M.;
7 Yu, K.; Chen, K.; Jiang, H.; Luo, C. Discovery of Novel Trimethoxy-Ring BRD4 Bromodomain
8 Inhibitors: AlphaScreen Assay, Crystallography and Cell-Based Assay †The Authors Declare No
9 Competing Interests. ‡Electronic Supplementary Information (ESI) Available. See DOI:
10 10.1039/C7md00083a. *Medchemcomm* **2017**, 8 (6), 1322–1331.
- 11 (39) Wang, K.; Kim, D.; Dömling, A. Cyanoacetamide MCR (III): Three-Component Gewald Reactions
12 Revisited. *J. Comb. Chem.* **2010**, 12 (1), 111–118.
- 13 (40) Mertz, J. A.; Conery, A. R.; Bryant, B. M.; Sandy, P.; Balasubramanian, S.; Mele, D. A.; Bergeron, L.;
14 Sims, R. J. Targeting MYC Dependence in Cancer by Inhibiting BET Bromodomains. *Proc Natl Acad*
15 *Sci U S A* **2011**, 108 (40), 16669–16674.
- 16 (41) Collignon, A.; Hospital, M. A.; Montersino, C.; Courtier, F.; Charbonnier, A.; Saillard, C.; D’Incan, E.;
17 Mohty, B.; Guille, A.; Adelaïde, J.; Carbuccioni, N.; Garnier, S.; Mozziconacci, M. J.; Zemmour, C.;
18 Pakradouni, J.; Restouin, A.; Castellano, R.; Chaffanet, M.; Birnbaum, D.; Collette, Y.; Vey, N. A
19 Chemogenomic Approach to Identify Personalized Therapy for Patients with Relapse or Refractory
20 Acute Myeloid Leukemia: Results of a Prospective Feasibility Study. *Blood Cancer J* **2020**, 10 (6), 64.
- 21 (42) Karjalainen, R.; Pemovska, T.; Popa, M.; Liu, M.; Javarappa, K. K.; Majumder, M. M.; Yadav, B.;
22 Tamborero, D.; Tang, J.; Bychkov, D.; Kontro, M.; Parsons, A.; Suvela, M.; Mayoral Safont, M.;
23 Porkka, K.; Aittokallio, T.; Kallioniemi, O.; McCormack, E.; Gjertsen, B. T.; Wennerberg, K.; Knowles,
24 J.; Heckman, C. A. JAK1/2 and BCL2 Inhibitors Synergize to Counteract Bone Marrow Stromal Cell-
25 Induced Protection of AML. *Blood* **2017**, 130 (6), 789–802.
- 26 (43) Benyamine, A.; Le Roy, A.; Mamessier, E.; Gertner-Dardenne, J.; Castanier, C.; Orlanducci, F.;
27 Pouyet, L.; Goubard, A.; Collette, Y.; Vey, N.; Scotet, E.; Castellano, R.; Olive, D. BTN3A Molecules
28 Considerably Improve Vγ9Vδ2T Cells-Based Immunotherapy in Acute Myeloid Leukemia.
29 *Oncoimmunology* **2016**, 5 (10), e1146843.
- 30 (44) Di Veroli, G. Y.; Fornari, C.; Wang, D.; Mollard, S.; Bramhall, J. L.; Richards, F. M.; Jodrell, D. I.
31 Combenefit: An Interactive Platform for the Analysis and Visualization of Drug Combinations.
32 *Bioinformatics* **2016**, 32 (18), 2866–2868.
- 33 (45) Bhadury, J.; Nilsson, L. M.; Muralidharan, S. V.; Green, L. C.; Li, Z.; Gesner, E. M.; Hansen, H. C.;
34 Keller, U. B.; McLure, K. G.; Nilsson, J. A. BET and HDAC Inhibitors Induce Similar Genes and
35 Biological Effects and Synergize to Kill in Myc-Induced Murine Lymphoma. *PNAS* **2014**, 111 (26),
36 E2721–E2730.
- 37 (46) Dickson, M. A.; Rathkopf, D. E.; Carvajal, R. D.; Grant, S.; Roberts, J. D.; Reid, J. M.; Ames, M. M.;
38 McGovern, R. M.; Lefkowitz, R. A.; Gonen, M.; Cane, L. M.; Dials, H. J.; Schwartz, G. K. A Phase I
39 Pharmacokinetic Study of Pulse-Dose Vorinostat with Flavopiridol in Solid Tumors. *Invest New*
40 *Drugs* **2011**, 29 (5), 1004–1012.
- 41 (47) Amorim, S.; Stathis, A.; Gleeson, M.; Iyengar, S.; Magarotto, V.; Leleu, X.; Morschhauser, F.; Karlin,
42 L.; Broussais, F.; Rezai, K.; Herait, P.; Kahatt, C.; Lokiec, F.; Salles, G.; Facon, T.; Palumbo, A.;
43 Cunningham, D.; Zucca, E.; Thieblemont, C. Bromodomain Inhibitor OTX015 in Patients with
44 Lymphoma or Multiple Myeloma: A Dose-Escalation, Open-Label, Pharmacokinetic, Phase 1 Study.
45 *Lancet Haematol* **2016**, 3 (4), e196–204.
- 46 (48) Berthon, C.; Raffoux, E.; Thomas, X.; Vey, N.; Gomez-Roca, C.; Yee, K.; Taussig, D. C.; Rezai, K.;
47 Roumier, C.; Herait, P.; Kahatt, C.; Quesnel, B.; Michallet, M.; Recher, C.; Lokiec, F.; Preudhomme,

- 1 C.; Dombret, H. Bromodomain Inhibitor OTX015 in Patients with Acute Leukaemia: A Dose-
2 Escalation, Phase 1 Study. *Lancet Haematol* **2016**, *3* (4), e186-195.
- 3 (49) Lewin, J.; Soria, J.-C.; Stathis, A.; Delord, J.-P.; Peters, S.; Awada, A.; Aftimos, P. G.; Bekradda, M.;
4 Rezai, K.; Zeng, Z.; Hussain, A.; Perez, S.; Siu, L. L.; Massard, C. Phase Ib Trial With Birabresib, a
5 Small-Molecule Inhibitor of Bromodomain and Extraterminal Proteins, in Patients With Selected
6 Advanced Solid Tumors. *J Clin Oncol* **2018**, *36* (30), 3007–3014.
- 7 (50) Zhao, L.; Okhovat, J.-P.; Hong, E. K.; Kim, Y. H.; Wood, G. S. Preclinical Studies Support Combined
8 Inhibition of BET Family Proteins and Histone Deacetylases as Epigenetic Therapy for Cutaneous T-
9 Cell Lymphoma. *Neoplasia* **2019**, *21* (1), 82–92.
- 10 (51) Romagnoli, R.; Prencipe, F.; Oliva, P.; Cacciari, B.; Balzarini, J.; Liekens, S.; Hamel, E.; Brancale, A.;
11 Ferla, S.; Manfredini, S.; Zurlo, M.; Finotti, A.; Gambari, R. Synthesis and Biological Evaluation of
12 New Antitubulin Agents Containing 2-(3',4',5'-Trimethoxyanilino)-3,6-Disubstituted-4,5,6,7-
13 Tetrahydrothieno[2,3-c]Pyridine Scaffold. *Molecules* **2020**, *25* (7), 1690.
- 14 (52) Cai, M.-G.; Wu, Y.; Chang, J. Synthesis and Biological Evaluation of 2-Arylimino-3-Pyridin-
15 Thiazolineone Derivatives as Antibacterial Agents. *Bioorg Med Chem Lett* **2016**, *26* (10), 2517–
16 2520.
- 17 (53) Filippakopoulos, P.; Knapp, S. The Bromodomain Interaction Module. *FEBS Letters* **2012**, *586* (17),
18 2692–2704.

