

1 **Low diversity and microdiversity of comammox bacteria in wastewater systems suggests**
2 **wastewater-specific adaptation within the *Ca. Nitrospira nitrosa* cluster.**

3

4 Irmario Cotto¹, Katherine J. Vilardi¹, Linxuan Huo², Emily C. Fogarty³, Wendell Khunjar⁴,
5 Christopher Wilson⁵, Haydee De Clippeleir⁶, Kevin Gilmore⁷, Erika Bailey⁸, Sebastian Lucker⁹,
6 Ameet J. Pinto^{2*}

7

8 1 Department of Civil and Environmental Engineering, Northeastern University, Boston, MA,
9 USA

10 2 School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA,
11 USA

12 3 Committee on Microbiology, The University of Chicago, Chicago, IL, USA

13 4 Hazen and Sawyer, Inc., New York, NY, USA

14 5 Hampton Roads Sanitation District, Norfolk, VA, USA

15 6 DC Water, Washington DC, USA

16 7 Department of Civil and Environmental Engineering, Bucknell University, Lewisburg, PA,
17 USA

18 8 City of Raleigh Public Utilities, Raleigh, NC, USA

19 9 Department of Microbiology, RIBES, Radboud University, Nijmegen, the Netherlands

20

21 *Corresponding author: ameet.pinto@ce.gatech.edu

22

23

24

25

26

27

28

29

30 **Abstract**

31 Studies have found *Ca. Nitrospira nitrosa*-like bacteria to be the principal or sole comammox
32 bacteria in nitrogen removal systems for wastewater treatment. In contrast, multiple populations
33 of strict ammonia and nitrite oxidizers co-exist in similar systems. This apparent lack of diversity
34 is surprising and could impact the feasibility of leveraging comammox bacteria for nitrogen
35 removal. We used full-length 16S rRNA gene sequencing and genome-resolved metagenomics to
36 compare population-level (i.e., species) diversity of comammox bacteria with that of strict
37 nitrifiers in full-scale wastewater treatment systems and assess whether these observations were
38 consistent or diverged at the strain-level. Full-length 16S rRNA gene sequencing indicated that
39 while *Nitrosomonas*-like bacteria exhibited higher population-level diversity, the effective
40 microdiversity of most *Nitrospira*-like bacteria were comparatively higher except for one
41 *Nitrospira* Lineage II population. Comammox bacterial metagenome assembled genomes (MAGs)
42 were associated with *Ca. Nitrospira nitrosa*. The average amino acid identity between comammox
43 bacterial MAGs ($93\% \pm 3$) across systems was significantly higher than that of the *Nitrosomonas*-
44 like ammonia oxidizers ($73\% \pm 8$) and the *Nitrospira*-like nitrite oxidizer MAGs ($75\% \pm 13$),
45 suggesting that the same comammox population was detected in all systems. Comammox bacteria
46 and some ammonia oxidizers MAGs were significantly less microdiverse than most ammonia and
47 nitrite oxidizers. Interestingly, strain-resolved analysis also indicates that different nitrogen
48 removal systems harbor different comammox bacterial strains within the *Ca. Nitrospira nitrosa*
49 cluster. These results suggest that comammox bacteria associated with *Ca. Nitrospira nitrosa* have
50 low species- and strain-level diversity in nitrogen removal systems and may thus harbor specific
51 adaptations to the wastewater ecosystem.

52

53 **Introduction**

54 Aerobic nitrification processes for nitrogen removal from wastewater are largely centered around
55 biotransformation and growth kinetics of strict ammonia oxidizing bacteria (AOB) and nitrite
56 oxidizing bacteria (NOB) (1). Since their discovery (2–5), several studies have detected
57 comammox bacteria in a wide range of environmental and engineered systems (2–9), including
58 nitrogen removal processes in laboratory and full-scale wastewater treatment bioreactors (9–20).
59 Compared to oligotrophic engineered systems such as drinking water (9,21) and even tertiary
60 treatment systems at full-scale wastewater treatment plants (WWTPs) (16) with low nitrogen
61 concentrations, most studies have identified *Candidatus Nitrospira nitrosa*-like comammox
62 bacteria as the principal or sole comammox bacteria in laboratory or full-scale secondary treatment
63 systems (11,22,23). This apparent lack of diversity of comammox bacteria in secondary treatment
64 processes is not only surprising but has the potential to impact treatment strategies centered around
65 comammox bacteria (24,25) as functional outcomes reliant on low diversity communities may be
66 less resilient to environmental fluctuations and perturbations.

67

68 *Nitrosomonas*-like AOB and *Nitrospira*-like NOB have been studied extensively in wastewater
69 treatment systems and studies often report the coexistence of multiple populations (26–28) likely
70 occupying different ecological niches (i.e., adapted to different conditions) within a complex
71 community. Functional redundancy of multiple co-existing populations can confer stability to
72 microbial ecosystems ensuring the long-term persistence (29,30). Despite the apparent low
73 population diversity of comammox bacteria compared to strict nitrifiers, they have exhibited
74 remarkable stability in secondary treatment systems (11,15,31). One plausible reason for their
75 temporal persistence could be that comammox bacteria may harbor intra-population (i.e., strain)

76 level diversity as compared to population level (i.e., species). Previous literature has demonstrated
77 that microdiversity within populations can not only allow them to adapt to rapidly changing
78 conditions, but also contribute to their persistence (7,32). Further, small variations in metabolic
79 capacity may underpin metabolic diversity and thus enable the co-existence of multiple closely
80 related strains within a single population. *Nitrospira*-like bacteria have been shown to harbor high
81 levels of microdiversity in drinking water (33,34) and wastewater (35) systems. Moreover, studies
82 have reported that closely related co-existing *Nitrospira* strains can exhibit differences in substrate
83 affinities, and utilization rates (35). Since, both levels of diversity (i.e., population and intra-
84 population) have been associated with microbial persistence and ecosystem stability, it is plausible
85 that the persistence of comammox bacteria may be associated with higher levels of microdiversity;
86 this would be consistent with observations for *Nitrospira*-like bacteria (35).

87

88 To test this hypothesis, we used a combination of full-length 16S rRNA gene sequencing and
89 hybrid assembly and binning approaches to systematically explore the population- and strain-level
90 diversity of nitrifiers in three full-scale nitrogen removal systems with different process
91 configurations. While 16S rRNA gene sequencing cannot provide information on genome content
92 or metabolic potential, the single nucleotide resolution across the full length 16S rRNA gene can
93 help determine differences in (micro)diversity among nitrifying populations (35,36). Nevertheless,
94 16S rRNA gene sequences cannot be used to reliably distinguish comammox bacteria from strict
95 *Nitrospira*-NOB. One approach to differentiate between the different guilds within *Nitrospira*
96 bacteria and to analyze their diversity at high resolution is to leverage a genome resolved approach
97 to obtain high quality metagenome assembled genomes (MAGs). Various methods exist to
98 quantify diversity (e.g., average nucleotide identity [ANI] and average amino acid identity [AAI])

99 and microdiversity (e.g., average nucleotide identity from reads and nucleotide diversity from
100 single nucleotide polymorphisms [SNPs]) within populations. Nevertheless, their applicability
101 depends on the quality of MAGs which can be impacted when relying on short read sequencing
102 (e.g., Illumina) in the presence of closely related strains (37); long-read sequencing on the
103 Nanopore platform can help mitigate this constraint (38,39). However, this requires high levels of
104 coverage to obtain polished consensus sequences with lower error rates compared to the raw data
105 (37). Here, we utilize a hybrid metagenomic assembly approach, including long- and short-read
106 data in an effort to assemble high quality MAGs for subsequent microdiversity analyses (37).

107

108 Thus, the overall objective of this study was to test the hypothesis that the widespread distribution
109 and persistence of nearly identical comammox bacterial populations in multiple secondary
110 wastewater treatment systems was due to the co-existence of multiple strains (i.e., high
111 microdiversity). To accomplish this, we used a combination of full-length 16S rRNA gene
112 sequencing and hybrid metagenome assembly and binning approach to systematically explore the
113 population and strain level diversity of nitrifiers in three full-scale nitrogen removal systems with
114 different process configurations. In doing so, we also aimed to assess factors that may influence
115 the inter- and intra-population diversity of co-existing nitrifiers.

116

117 **2.0 Materials and Methods**

118 **2.1 Samples selection and processing**

119 Samples were selected from three nitrogen removal systems with different process configurations
120 (i.e., sequencing batch reactor (SBR), Integrated Fixed Film Activated Sludge (IFAS), and
121 Bardenpho 4-stages activated sludge system) with high concentrations of comammox bacteria in

122 our previous study (31). From this sample archive collected in 2017-2018, a total of thirty-three,
 123 twenty, and six samples, respectively, were selected for full-length 16S rRNA gene sequencing on
 124 the PacBio Sequel IIe platform, and short and long read metagenomic sequencing on the Illumina
 125 NovaSeq and Nanopore MinION platforms, respectively (Table 1). Sample collection and
 126 processing (including DNA extraction) and process data collection were described previously (31)
 127 and are outlined in the supplementary text.

128 **Table 1: Overview of system process type and sub-type, operational scale and samples included in**
 129 **this study and sample-specific sequencing strategy.**

<i>Site code</i>	<i>Process type</i>	<i>Process sub-type</i>	<i>Treatment stream</i>	<i>Selected samples (sampling month/sampling year)</i>
GRE	ND ¹	SBR ²	Mainstream	06/17 ⁺ , 07/17 ^{*†} , 08/17 ⁺ , 09/17 ^{*†} , 10/17 ^{*†} , 11/17 ⁺ , 12/17 ^{*†} , 02/18 ^{*†} , 03/17 ⁺ , 04/18 ^{*†} , 05/18 ⁺ , 06/18 ^{*†}
JAMMSM	ND ¹	IFAS ³	Mainstream	07/17 ⁺ , 09/17 ^{*†} , 10/17 ⁺ , 11/17 ^{*†} , 12/17 ⁺ , 01/18 ^{*†} , 02/18 ⁺ , 03/18 ^{*†} , 04/18 ^{*†} , 06/18 ^{*†}
NEU	ND ¹	Four/five stage	Mainstream	06/17 ⁺ , 07/17 ^{*†} , 08/17 ⁺ , 09/17 ^{*†} , 10/17 ^{*†} , 11/17 ^{*†} , 12/17 ⁺ , 01/18 ^{*†} , 02/17 ⁺ , 03/18 ^{*†} , 04/17 ⁺ , 05/18 [*] , 06/18 ⁺

130 ¹Nitrification-denitrification, ²Sequencing batch reactor, ³Integrated Fixed-film Activated Sludge

131 *Samples selected for Illumina NovaSeq sequencing

132 †Samples selected for PacBio full-length 16S rRNA gene sequencing

133 ‡Samples selected for nanopore sequencing on the MinION platform

134
135

136 **2.2 PacBio full-length 16S rRNA gene sequencing and data analysis**

137 Sample DNA extracts (Table 1) were sent for full-length 16S rRNA gene sequencing at the Roy J.
 138 Carver Biotechnology Center (Sequencing Core, University of Illinois Urbana-Champaign). The
 139 full-length 16S rRNA gene amplicons were generated with universal barcoded primers (27F and
 140 1492R)(40), PCR products were subject to library preparation and sequenced on the PacBio Sequel
 141 IIe using the circular consensus sequencing (CCS) mode. Raw reads were demultiplexed and CCS
 142 analyses were performed to obtain consensus reads. Further details on PCR, library preparation,
 143 and sequencing are provided in the supplementary text. Downstream data processing was
 144 performed on a system-by-system basis using DADA2 v1.19.2 (41) in R v4.0.4. using the sample
 145 inference method for full-length 16S rRNA gene with single-nucleotide resolution (42). Table S1a

146 summarizes the reads per samples at different stages of data processing in DADA2. ASVs with
147 ≥ 100 reads in any of the three systems ($\sim > 0.01\%$ relative abundance) were clustered into
148 operational taxonomy units (OTUs) using the function ‘IdClusters’ in DECIPHER v2.16.1. (43)
149 at 98.7% sequence identity, a previously recommended threshold for clustering full-length 16S
150 rRNA gene sequences at the species level (44,45). The most abundant ASV within each OTU was
151 used as representative sequence and its taxonomic affiliation was used as the consensus taxonomy
152 for the OTU. Additional details on ASV to OTU data processing are provided in the supplementary
153 text. Representative ASV sequence from all OTUs were aligned with MUSCLE v3.8.1551 (46),
154 and the phylogenetic tree was constructed and visualized using IQ-TREE v2.0.3 (47) and iTOL
155 v2.1.7 (48), respectively. Principal Coordinates Analysis (PCoA) was performed with the weighted
156 and unweighted UniFrac distance metric (49) using the ‘ordinate’ function of phyloseq v1.32.0
157 and the ‘plot_ordination’ function of ggplot2 v3.3.5 (50).

158

159 ASVs classified at the genus level as *Nitrospira* and *Nitrosomonas* (the only two nitrifying genera
160 detected in this study) were extracted from the rarefied ASV table (rarefied to the sample with
161 lowest read count) and reference 16S rRNA gene sequences for the two genera were obtained from
162 the SILVA SSU 138.1 database (Tables S2 and S3). Comammox bacterial 16S rRNA genes were
163 extracted from references genomes downloaded from NCBI (Table S4). MUSCLE v3.8.1551 was
164 used to align ASVs sequences from each genus with their respective references, and a maximum
165 likelihood phylogenetic tree of each species was generated using IQ-TREE v2.0.3 and visualized
166 in iTOL v2.1.7. To assess the population diversity (i.e., genus level) the Shannon diversity was
167 calculated per sample from the relative abundance of the OTUs classified as the same genus (e.g.,
168 *Nitrospira*) as the exponential of the Shannon index. To estimate intra-population diversity, the

169 effective microdiversity of each OTU was calculated on a per sample basis from the relative
170 abundance of its ASVs as the exponential of the Shannon index, which is analogous to the effective
171 number of strains (i.e., ASVs) within a population (i.e., OTU) (36).

172

173 **2.3 Illumina and Oxford Nanopore sequencing, co-assembly and hybrid assembly**

174 Seven samples from GRE and NEU and six samples from JAMMSM were sent for sequencing on
175 the Illumina NovaSeq 6000 platform, and two samples from each system on the Oxford Nanopore
176 Technologies GridIONx5 (Table 1) to the Roy J. Carver Biotechnology Center at the University
177 of Illinois Urbana-Champaign Sequencing Core. Library preparation and sequencing details are
178 provided in the supplementary text. These runs resulted in 1.78 billion paired-end short reads
179 (2×150 nt reads) and 45.3 Gbps of long read data (Table S1b and c). Raw short reads were filtered
180 using fastp v0.20.0 (51) and the Univec database was used to remove contamination from the
181 filtered reads as previously described (31). The resulting bam files were sorted using SAMtools
182 v1.9 (52) and converted into fastq files with bedtools v2.29.0 (53). All reads from the same system
183 were co-assembled into contigs using metaSpades v3.13.0 (54). Co-assemblies were performed
184 with kmer sizes of 21, 33, 55 and 77. Contigs smaller than 500 bp were removed from the co-
185 assemblies using the Anvi'o v6.1 command 'anvi-script-reformat-fasta' (55). The reformatted
186 assembly fasta files were indexed with bwa index v0.7.17 (56) and the paired end reads from each
187 metagenome were mapped to the respective co-assembly using bwa mem v0.7.17. The resulting
188 sam files were converted to bam files using 'samtools view -F 4 -bhS' to retain only mapped reads.
189 Hybrid metagenomic assemblies were performed using OPERA-MS v0.9.0., which combines the
190 advantages of short and long-read technologies to improve genome assemblies (37). The OPERA-
191 MS inputs were short-read metagenomic assemblies to provide a good representation of the

192 sequences in the metagenome, and long and short reads to identify connections between the contigs
193 and obtain contiguous assemblies with low base-pair error. OPERA-MS was executed with the
194 flags `--no-ref-clustering` and `--long-read-mapper minimap2`. Both sets of co-assemblies were
195 evaluated using QUAST v5.0.2 (57) (Table S5).

196

197 **2.4 Recovery, annotation, refining and dereplication of metagenome assembled genomes** 198 **(MAGs)**

199 Binning was performed separately, with co-assemblies from Illumina short reads only and hybrid
200 assemblies from each system using MetaBAT2 v2.12.1 (58), CONCOCT v1.1.0 (59), and
201 MaxBin2 v2.2.7 (60) using contigs greater than 2000 bp. Pilon v1.23 (61) was used for polishing
202 to improve the draft bins obtained from the hybrid assemblies. The quality and taxonomy of the
203 resulting bins were determined with CheckM v1.1.2 (62) and the Genome Taxonomy Database
204 Toolkit (GTDB-Tk 1.1.1, database release r86 v3) (63), respectively. Bins were subject to gene
205 calling using Prodigal v2.6.3 (64) and gene annotation against the KEGG database (65) using
206 kofamscan v1.2.0 (66). Only bins that were taxonomically assigned to known nitrifying genera or
207 those containing genes associated with nitrification (i.e., *amoA* [KO number K10944], *amoB*
208 [K10945], *amoC* [K10946], *hao* [K10535], *nxrA* [K00370], *nxrB* [K00371]) were retained for
209 manual refinement with Anvi'o v6.1. DASTool v1.1.2 (67) was used to combine and curate the
210 refined bins from the three binning methods and generate a non-redundant set of bins from each
211 co-assembly (i.e., one set of bins per system for the short-read only and for the hybrid assemblies).
212 In total, 43 nitrifiers bins from the short-read co-assemblies and 30 from the hybrid assemblies
213 were obtained. These bins were de-replicated using drep v2.5.4 at 95% ANI with completeness
214 and contamination thresholds set to 50% and 10%, respectively. This resulted in 44 nitrifier MAGs

215 with completeness and redundancy estimates higher than 50% and lower than 10%, respectively,
216 of which 21 contained genes associated with nitrification (i.e., *amoA*, *amoB*, *amoC*, *hao*, *nxrA*,
217 and/or *nxrB*).

218

219 **2.5 Taxonomy aware re-assembly using nitrifying bacterial MAGs**

220 We performed re-assemblies from reads mapped to the 44 nitrifying bacterial MAGs to improve
221 MAG quality. First, contigs associated with *Nitrospira* (including *comammox*) and *Nitrosomonas*
222 MAGs from each system (i.e., GRE, JAMMSM, NEU) were collated into a single fasta file per
223 system. The resulting fasta files were indexed using *bwa index v0.7.17* and short reads from the
224 respective system were mapped using *bwa mem v0.7.17*. The resultant bam file was split into
225 *Nitrospira* and *Nitrosomonas* specific bam files per system and respective fastq files were
226 generated per using *samtools fastq*. Long reads were also recruited by mapping the reads from
227 Nanopore sequencing to the fasta files with ‘*bwa mem -x ont2d*’. Short reads mapping to the
228 contigs of the MAGs classified as *Nitrospira* and *Nitrosomonas* were re-assembled using
229 *metaSpades v3.13.0* with *kmers 21, 33, 55 and 77* on a system-by-system basis. The new genus-
230 specific assemblies and fastq files of mapped long reads were used as input for OPERA-MS to
231 perform hybrid metagenomic re-assemblies (n=9). Quality assessment, binning, taxonomy
232 annotation, manual refining and gene calling were performed as described before. This resulted
233 in 36 nitrifier MAGs with completeness and redundancy estimates higher than 50% and lower than
234 10%, respectively, of which 24 contained genes associated with nitrification (*amoA*, *amoB*, *amoC*,
235 *hao*, *nxrA*, and/or *nxrB*). The abundance of each MAG per sample (reads per kilobase million,
236 RPKM) and proportion of genome covered was calculated with *coverM (68)*. Four low abundant
237 MAGs (3 AOB and 1 *Nitrospira*-NOB) with less than 50% genome coverage in any of the samples

238 were removed from subsequent analyses. The entire workflow for assembly and re-assembly of
239 MAGs is outlined in Figure S1. The nitrifier MAGs were subsequently phylogenetically placed in
240 the context of reference *Nitrospira* (Table S4) and *Nitrosomonas* (Table S6) genomes
241 (supplementary text).

242

243 **2.6 Diversity and micro-diversity analysis of nitrifiers**

244 FastANI v1.3 (69) was used to calculate the pairwise average nucleotide identity (ANI) between
245 MAGs within each functional group (i.e., AOB, *Nitrospira*-NOB, and *Nitrospira*-comammox).
246 ANI represents the mean nucleotide identity of the orthologous genes shared between two genomes
247 offering a robust resolution between similar or identical species (i.e., ~80-100% ANI) (69). Since
248 ANI values lower than 80% are not reported, we also calculated the average amino acid identity
249 (AAI) to estimate similarity between two genomes at the amino acid level using compareM (70).
250 To determine intra-population diversity (i.e., microdiversity), we calculated the average nucleotide
251 identity from reads (ANIr) (32) with 90% read identity threshold, as recommended for intra-
252 population comparisons. inStrain v1.3.9 (71) was used to determine the nucleotide diversity for
253 each MAG in each sample and their population average nucleotide identity (popANI) between the
254 samples where they were detected. Nucleotide diversity is a measurement of genetic
255 (micro)diversity at every position along the genome using mapped reads, while popANI is a unique
256 ANI calculation performed by inStrain that considers both major and minor alleles. This is
257 different from the traditional ANI (called consensus ANI [conANI] in inStrain), which only
258 considers major alleles to call (or not) a substitution. Details on parameters and procedures
259 associated with the implementation of ANIr and inStrain are presented in the supplementary text.
260

261 **2.7 Statistical Analyses**

262 Statistical tests were performed using R v4.0.4. All correlations were performed using a linear
263 regression model. The Spearman's rank correlation coefficients between MAG abundances were
264 performed with the 'rcorr' function of the R package Hmisc v4.5.0. Pair-wise significances were
265 calculated with the Kruskal-Wallis test, a nonparametric approach to the one-ANOVA. Principal
266 coordinates analysis (PCoA) with the weighted and unweighted UniFrac distance metrics was used
267 to compare the community composition among systems.

268

269 **3.0 Results**

270 **3.1 Community composition in nitrogen removal systems**

271 Three nitrification-denitrification systems with high absolute abundance of comammox bacteria
272 were selected for this time-series study on the basis of our previous findings (31) since high
273 sequencing coverage is critical for evaluation of sub-population level diversity. Specifically, over
274 the sampling period (June 2017 to June 2018) comammox bacteria constituted approximately 0.77
275 (± 0.32), 4.7 (± 4.21), and 0.45 (± 0.24) % of total bacterial 16S rRNA gene abundances at GRE,
276 JAMMSM, and NEU, respectively.


277

278 The PacBio full-length 16S rRNA gene sequencing resulted in 2,317,019 total reads (Table S1a)
279 with 7501, 5783, and 8260 ASVs at GRE, JAMMSM and NEU, respectively. Each ASV set was
280 rarefied to the sample with the smallest library size per system resulting in 7040, 4349, and 5749
281 ASVs from GRE, JAMMSM, and NEU, respectively, and a total of 16651 unique ASVs across all
282 three systems. ASVs with a total of 100 reads in each system were clustered into OTUs at 97%
283 identity, resulting in 846 OTUs (Figure 1A; Table S7). The most abundant OTUs at GRE and NEU

284 were from the classes *Gammaproteobacteria* and *Alphaproteobacteria* and the phylum
285 *Bacteroidota* while the class *Nitrospira* had the highest relative abundance in JAMMSM. PCoA
286 using weighted (WUF) and unweighted UniFrac (UUF) distance metrics (Figure 1B) demonstrated
287 that samples clustered by system and the community structure between GRE and NEU (WUF:
288 0.18 ± 0.03 ; UUF: 0.46 ± 0.03) were significantly more similar than GRE and JAMMSM (WUF:
289 0.28 ± 0.02 ; UUF: 0.56 ± 0.05) and NEU and JAMMSM (WUF: 0.29 ± 0.03 ; UUF: 0.52 ± 0.04)
290 ($p < 0.05$; Figure S2A and B). The greater similarity between GRE and NEU is likely due to the
291 fact that these were suspended phase communities as compared to attached phase (i.e., biofilm
292 samples) communities collected from the IFAS system at JAMMSM.

293


294 *Nitrosomonas*- (31, 25, and 14 ASVs at GRE, JAMMSM, and NEU, respectively) and *Nitrospira*-
295 like bacteria (37, 36, and 29 ASVs at GRE, JAMMSM, and NEU, respectively) were the only
296 nitrifiers present in the systems. *Nitrospira* lineage I ASVs were the most abundant group across
297 all systems with average relative abundances of 2.91, 25.13, and 2.88%, followed by *Nitrospira*
298 lineage II with average relative abundances 1.41, 7.07, and 0.64% in GRE, JAMMSM, and NEU,
299 respectively (Figure 1C). *N. oligotropha* (Cluster 6a) and *N. communis* (Cluster 8) -like ASV's
300 were present in all systems with relative abundances of 0.5 and 0.33, 1.68 and 0.93, and 0.58 and
301 0.19% in GRE, JAMMSM, and NEU, respectively. *N. europaea/mobilis* (Cluster 7)-like ASVs
302 were also detected in GRE and JAMMSM at very low abundances compared to the other two
303 *Nitrosomonas* lineages (average ~ 0.032 and 0.002%, respectively).



304
 305 **Figure 1:** Community composition of the nitrogen removal systems. (A) Phylogenetic tree constructed using full-length 16S rRNA
 306 gene sequences of the representative ASV in each OTU (98.7% sequence similarity cluster), with their corresponding relative
 307 abundances shown at the right. Branches are collapsed at the phylum/class level and corresponding number of OTUs within each
 308 phylum/class across all three systems are indicated. Black circles on branches designate bootstrap support. Bubbles represent
 309 cumulative relative abundances of all OTUs within each phylum/class per system. (B) Principal Coordinate Analysis of community
 310 composition with unweighted and weighted UniFrac distance metrics using OTU data. Points represent the samples. (C)
 311 Cumulative relative abundances of OTUs corresponding to *Nitrospirira* Lineages and *Nitrosomonas* clusters per system at each
 312 time point analyzed. Colors correspond to each nitrogen removal system with (GRE, pink; JAMMSM, green; NEU, blue).

313 3.2 16S rRNA gene sequence-based diversity and microdiversity of nitrifiers

314 The *Nitrospira*-like ASVs belonged to lineage I (20, 24, and 22 ASVs at GRE, JAMMSM, and
315 NEU, respectively) or lineage II (17, 12, and 7 at GRE, JAMMSM, and NEU, respectively) (Figure
316 S3A), while *Nitrosomonas*-like ASVs were associated with *N. oligotropha* (23, 21, and 10 at GRE,
317 JAMMSM, and NEU, respectively), *N. europaea/mobilis* (5 and 1 at GRE and JAMMSM,
318 respectively), and *N. communis* (3, 3, and 4 at GRE, JAMMSM, and NEU, respectively) lineages
319 (Figure S3B). The *Nitrospira*- and *Nitrosomonas*-like ASVs clustered into 3, 3 and 2, and 21, 14
320 and 10 OTUs for GRE, JAMMSM and NEU, respectively. The Shannon diversity (i.e., population
321 diversity) of *Nitrosomonas*-like OTUs was significantly higher than *Nitrospira*-like OTUs, while
322 the effective microdiversity of *Nitrospira*-like OTUs was significantly higher than *Nitrosomonas*-
323 like OTUs in all systems ($p < 0.05$; Figure 2A,). *Nitrospira* lineage I OTUs consistently showed the
324 highest microdiversity (Figure 2B) while two *Nitrospira* lineage II OTUs from GRE and
325 JAMMSM had a lower effective microdiversity than most nitrifier OTUs. These results suggest
326 high population level diversity and low microdiversity for *Nitrosomonas*-like bacteria, while the
327 opposite was observed for *Nitrospira*-like bacteria in Lineage I (more ASVs per OTU). The
328 effective microdiversity of *Nitrospira*-like OTUs was positively correlated with their relative
329 abundance (Figure S4A). However, this was neither the case for *Nitrosomonas*-like OTUs (Figure
330 S4B) and the overall bacterial community (Figure S4C). Although the relative abundances of some
331 other highly microdiverse OTUs were positively correlated with effective microdiversity (e.g.,
332 *Hyphomicrobium*), this correlation was not statistically significant for most (e.g., Figure S5). This
333 suggests that the increase in effective microdiversity of *Nitrospira*-like OTUs with increasing
334 relative abundance is more likely to be an ecological phenomenon rather than a sampling artifact.



335

336 **Figure 2:** Diversity and microdiversity based on ASV analyses. (A) Boxplot of Shannon diversity and effective microdiversity of
 337 *Nitrospira* (lighter shade) and *Nitrosomonas* (darker shade) at each system. (B) Average effective microdiversity of major OTUs
 338 per system. *Nitrospira* and *Nitrosomonas* OTUs are colored by system (GRE, pink; JAMMSM, green; NEU, blue), represented by
 339 closed and open symbols, respectively.

340

341

342

343

344

345 **3.3 Improving the recovery of metagenome assembled genomes of nitrifying bacteria**

346 A total of 43 dereplicated nitrifiers MAGS were recovered from short-read assemblies. However,
347 several MAGs lacked key genes associated with nitrification (i.e., *amoA*, *amoB*, *amoC*, *hao*, *nxrA*,
348 *nxB*). For example, only 2 of the 7 *Nitrosomonas* MAGs (from JAMMSM and NEU) contained
349 the complete *amoCAB* operon and only one had the *hao* gene. To improve MAG quality, we
350 incorporated long reads from Nanopore sequencing and a hybrid assembly using OPERA-MS (37).
351 The hybrid approach increased the contiguity of the assembly and resulted in 30 nitrifier MAGs,
352 with nitrification genes present in 19 MAGs (Table S8). It resulted in significant improvement of
353 MAGs retrieved from JAMMSM but not for GRE or NEU. Thus, we pursued taxonomy aware re-
354 assembly (72) of the 44 non-redundant nitrifier MAGs after dereplicating bins from short-read and
355 hybrid assembly approach. The resultant 36 nitrifier MAGs had completeness greater than 70%
356 and redundancy lower than 10% (Table S8), with nitrification genes present in 24 MAGs.
357 Although the number of retrieved nitrifier MAGs was lower after the reassembly process, there
358 was significant improvement in quality, especially for comammox and AOB genomes. For
359 example, both analyses (before and after reassembly) resulted in 4 *Nitrospira*-comammox MAGs.
360 However, 2 of the 4 comammox MAGs recovered after reassembly contained the entire genetic
361 repertoire required for nitrification, and the other two lack only one gene. In contrast, all pre-
362 reassembly comammox MAGs lacked at least one nitrification gene. These improvements were
363 also observed for AOB including significant improvement in MAG quality (i.e., higher
364 completeness, lower redundancy, and less fragmentation) (Table S8). These improvements
365 resulted from longer contigs created during the reassembly process compared with previous co-
366 assemblies. In contrast to comammox and AOB, despite becoming less fragmented post-
367 reassembly, fewer *Nitrospira*-NOB MAGs contained all genes associated with nitrite oxidation.


368 3.4 Phylogenomic placement and abundance of nitrifier MAGs

369 The 11 canonical NOB and 4 comammox were associated with *Nitrospira* lineage I (8 *Nitrospira*-
370 NOB) and *Nitrospira* lineage II (3 *Nitrospira*-NOB and 4 *Nitrospira*-comammox), with all
371 comammox MAGs closely related with *Ca. Nitrospira nitrosa* (Figure 3A) and belonged to
372 *Nitrospira*-comammox clade A.1 based on *hao* gene phylogenetic inference (Figure S6). This is
373 in agreement with a previous study that suggests comammox clade A.1 usually co-occur with the
374 *Nitrospira* lineage I populations in wastewater treatment plants (WWTPs) while clade A.2 and
375 most clade B are typically found in drinking water treatment plants (DWTPs) (21). While non-
376 WWTP settings (e.g., soils, sediments, lakes, rivers, DWTPs) harbor diverse comammox bacteria
377 (9,18,21,73,74), our findings suggest very low diversity of comammox bacteria in wastewater
378 systems which is consistent with other wastewater studies (12,15,17,22,75–77). All AOB MAGs
379 (n=20) were associated with the *N. oligotropha* (n=14) and *N. communis* (n=3) lineages (Figure
380 3B). Although several members of the *N. europaea/mobilis* lineage have been detected in sewage
381 treatment plants (78), only three AOB MAGs related to this lineage were recovered from
382 JAMMSM at very low abundances and less than 50% genome coverage. Thus, these MAGs were
383 excluded from subsequent analyses. The three *N. communis* cluster MAGs (GRE AOB 1,
384 JAMMSM AOB 4, NEU AOB 3) were nearly 100% identical but shared less than 75% ANI with
385 all reference genomes. This lineage can be divided into the *N. communis* and the *N. nitrosa* clusters
386 (79), with *N. communis* species urease negative and detected primarily in agricultural soils, and
387 isolates of *N. nitrosa* urease positive, preferring aquatic habitats and often found in wastewater
388 treatment plants (80). Our three MAGs were urease positive but still distant to *N. nitrosa* in the
389 phylogenomic analysis (Figure 3B), suggesting the presence of a novel phylogenetic lineage.
390 Lastly, most species of the *N. oligotropha* lineage have been recovered from oligotrophic

391 freshwaters and almost all are urease positive (78,80). Accordingly, 12 of the 14 AOB MAGs
392 placed within this *Nitrosomonas* lineage were urease positive.

393

394 *Nitrospira*-NOB was the most abundant functional group in all three systems for the duration of
395 the study with higher abundances in JAMMSM (314.5 ± 70.6 RPKM) as compared to NEU (178.2
396 ± 12.5 RPKM) and GRE (149.3 ± 11.7 RPKM). *Nitrosomonas*-like MAGs were more abundant in
397 NEU (105.2 ± 23 RPKM) and GRE (86 ± 15.8 RPKM) as compared to JAMMSM (53.2 ± 27.4).
398 This suggests that *Nitrospira*-like bacteria may prefer biofilm-based growth over suspended phase
399 (81). *Nitrospira*-comammox relative abundances were very similar in all systems, ranging from
400 55.5 to 64.9 RPKM in JAMMSM and NEU, respectively. The abundances of nitrifying MAGs
401 were weakly correlated with each other, with only 37 of 195 correlations significant (p-value
402 < 0.05). Most of the significant correlations between AOB and *Nitrospira*-NOB were positive
403 (Figure S7), likely arising as consequence of metabolic interactions of these groups within the
404 nitrification process. Comammox bacteria had very few correlations with other nitrifiers, with
405 primarily negative correlations with MAGs from *N. oligotropha*-like bacteria (cluster 6a).




406

407 **Figure 3:** Phylogenetic placement of (A) *Nitrospira* and (B) *Nitrosomonas* MAGs (GRE, pink; JAMMSM, green and NEU, blue)
 408 with 78 and 48 reference genomes (black), respectively. Comammox clade A (black arrow), clade B (grey arrow) and canonical
 409 NOB (white arrow) are indicated by arrows and phylogenetic lineages by colored boxes. Branches with reference genomes
 410 that did not include MAGs from this study were collapsed and labeled with a representative reference genome. All reference genomes
 411 used for the reconstruction of the trees are listed in Tables S4 and S6. (C) Cumulative abundances (RPKM) of *Nitrospira*-
 412 comammox, *Nitrospira*-NOB and *Nitrosomonas* MAGs at each time point per system.

413

414 3.5 Population-level diversity of nitrifying bacteria within and between systems

415 Post dereplication at 95% ANI (69,82), pairwise ANI and AAI were calculated between all MAGs
416 within the same functional group (i.e., AOB, *Nitrospira*-NOB, and *Nitrospira*-comammox)
417 (Figure 4A-B, Tables S9a-b). Of the 136 pairwise comparisons between the 17 AOB MAGs, only
418 18 (13.2%) had ANI values higher than 80% suggesting high population-level diversity within and
419 between nitrogen removal systems for the *Nitrosomonas*-like AOB (Figure 4A). The pairwise AAI
420 for *Nitrospira*-NOB MAGs showed a bimodal distribution (Figure 4B) reflective of the two
421 lineages detected in each system (i.e., Lineage I and II). Pairwise AAIs for *Nitrospira*-NOB MAGs
422 of the same lineage within or across systems were typically >85% (Table S9b). These results
423 suggest that while multiple AOB and *Nitrospira*-NOB coexist in each system, AOB were
424 substantially more diverse than *Nitrospira*-NOB. In contrast, three of the four comammox MAGs
425 (one from each system) share ANI values between 95-97% and are closely related with *Ca.*
426 *Nitrospira nitrosa* (31). Although a comammox MAG from JAMMSM (JAMMSM CMX 2) has
427 lower than 95% with the other three MAGs, it still exhibited greater than 85% AAI (Figure 4B)
428 with them and also falls within the *Ca. Nitrospira nitrosa* cluster (Figure 3A)(83). Pairwise AAI
429 comparisons (which are highly correlated with ANI – Figure S8) indicated that *Nitrospira*-
430 comammox MAGs have significantly higher AAI compared with AOB and NOB, while AOB
431 were the most diverse functional group at the population level ($p < 0.05$; Figure 4B). Moreover,
432 similar to ASV analysis, *Nitrospira*-NOB MAGs were associated with one of two clusters
433 (*Nitrospira* lineage I or II), with highly similar populations within each cluster (Figure 4B) being
434 detected across systems. These findings further confirm that in contrast to AOB that demonstrate
435 high population diversity, comammox bacteria show very low population diversity and, in fact,
436 may all be associated with the same population across multiple wastewater systems



437

438 **Figure 4:** (A) ANI values calculated between *Nitrosomonas* (AOB), *Nitrospira*-NOB and *Nitrospira*-comammox (CMX) MAGs along
 439 with the phylogenetic placement of the MAGs. MAG labels are colored by system (GRE, pink; JAMMSM, green and NEU, blue).
 440 (B) AAI values per functional group. Black lines represent the average, points represent the AAI value between two MAGs and
 441 colors decode the systems that are compared. Pairwise statistical comparisons were performed with the Kruskal-Wallis test.

442

443


444 3.6 . Microdiversity of nitrifying bacteria

445 The ANIr distributions for comammox and AOB MAGs were not significantly different from each
446 other, and significantly higher than for *Nitrospira*-NOB MAGs (Figure 5A). While *Nitrospira*-
447 NOB overall demonstrated lower ANIr indicating higher microdiversity compared to AOB and
448 comammox bacteria, there was some variability within each system (Figure S9). These observed
449 trends were similar between ANIr and inStrain estimated nucleotide diversity (Figure 5B), with
450 AOB showing on average lowest microdiversity and *Nitrospira*-NOB the highest. Specifically, the
451 average nucleotide diversity of *Nitrosomonas*-like AOB was significantly lower than of both
452 *Nitrospira* groups, i.e., NOB and comammox ($p < 0.05$), while with ANIr only *Nitrospira*-NOB
453 were significantly more microdiverse than the two other functional groups. This difference arises
454 from the fact that the two measures of intra-population diversity (i.e., ANIr and nucleotide
455 diversity) are calculated differently. For instance, ANIr considers only the major alleles in the
456 consensus sequence to call a substitution, while nucleotide diversity is calculated using base pair
457 frequencies at each position. Nonetheless, both approaches indicate that *Nitrospira*-NOB exhibit
458 significantly higher microdiversity as compared to *Nitrosomonas*-like AOB and *Nitrospira*-
459 comammox bacteria. These results are consistent with the 16S rRNA gene analysis where
460 *Nitrospira* lineage I and most lineage II OTUs had significantly higher effective microdiversity
461 than the *Nitrosomonas*-like OTUs and the remaining *Nitrospira* lineage II OTUs. Based on these
462 results, we speculate that the *Nitrospira* lineage II OTUs with low effective microdiversity likely
463 correspond to comammox bacteria (Figure 2B).

464

465 Finally, popANI, a microdiversity-aware ANI calculation, was used to discriminate between
466 strains across samples using the recommended popANI threshold of 99.999% (Olm et al., 2021).

467 All popANI values from GRE_CMX_1, JAMMSM_CMX_1, and NEU_CMX_1 were above or
468 very close to the 99.999% threshold, suggesting low strain-level diversity of comammox bacteria
469 (Figure 5C). Only one comammox genome retrieved from JAMMSM (JAMMSM_CMX_2) had
470 popANI values lower than the recommended threshold. However, the average relative abundance
471 of this comammox (RPKM = 7 ± 2) was particularly low as compared to the more abundant
472 comammox in JAMMSM (RPKM = 48 ± 21) and those retrieved from GRE (RPKM = 60 ± 19) and
473 NEU (RPKM = 65 ± 17). Therefore, the principal or sole comammox bacteria in each system is
474 not only a single population but likely also a specific strain. The ANI_r for each comammox MAG
475 estimated by mapping reads from a different system (e.g., mapping GRE or NEU reads to MAGs
476 assembled from JAMMSM) were lower than when mapping of reads from the system from where
477 the MAG was obtained (e.g., mapping JAMMSM reads to MAGs assembled from JAMMSM;
478 Figure S10). This suggests that on the one hand the comammox bacterial population within a single
479 system is restricted to a specific *Ca. Nitrospira nitrosa*-like strain, but on the other hand that
480 different systems contain different strains.



481

482 **Figure 5:** (A) ANI and (B) nucleotide diversity values from each functional group. Points represent the average value of each MAG
 483 and are colored according to the systems (GRE, pink; JAMMSM, green and NEU, blue) from which the MAGs were recovered. (C)
 484 popANI values from each MAG. MAGs are colored by system and points represent the popANI value of the MAG between two
 485 samples.

486

487

488

489

490

491

492

493

494

495 4 Discussion

496 4.1 Hybrid assembly and re-assembly processes improve the quality of metagenome

497 assembled genomes

498 Reliance on short reads for genome assembly from complex metagenomes can lead to incomplete
499 and fragmented MAGs especially in systems containing closely related populations (84). Indeed,
500 short read analysis resulted in highly fragmented nitrifier MAGs lacking several nitrification
501 genes. To improve the short read assemblies, we used long reads generated from Nanopore
502 sequencing with a hybrid assembly approach (38). Recent studies have demonstrated the ability of
503 hybrid assemblies to significantly improve the recovery of complex bacterial genomes (85,86) by
504 increasing the contiguity and accuracy of metagenome assembly (87). Combining the two
505 sequencing techniques (i.e., short and long reads) with the hybrid assembler OPERA-MS resulted
506 in modest assembly and binning improvements regarding genome contiguity and completeness.
507 However, the bead beating based DNA extraction protocol may have resulted in shearing of
508 genomic DNA into smaller fragments which is not ideal for long-read sequencing (88) and resulted
509 in lower read lengths. In an attempt to further improve MAG quality, we performed taxonomy
510 aware reassemblies (72) as described in the materials and methods section. Since a hybrid approach
511 resulted in some improvements in MAG quality, we also adopted a hybrid strategy with OPERA-
512 MS for taxonomy aware reassembly. This significantly improved the completeness and contiguity
513 of MAGs and number of annotated nitrifier genes (especially for *Nitrosomonas* and comammox
514 *Nitrospira*) binned within them. While the *Nitrospira*-NOB MAGs also became less fragmented,
515 they showed fewer improvements with respect to the presence of nitrite oxidation genes. This is
516 likely caused by the high levels of microdiversity within *Nitrospira*-NOB (Figures 2 and 5), which
517 likely impacts both the read mapping and de novo assembly process. In the future, this issue might

518 be improved by incorporating more long-read data into the re-assemblies, especially once the
519 accuracy of long-read sequencing technologies further increases.

520

521 **4.2 *Ca.* Nitrospira nitrosa-like bacteria are the primary comammox bacteria in secondary**
522 **wastewater treatment for nitrogen removal**

523 Structural diversity and functional redundancy are inherently linked to process stability when
524 environmental and process conditions vary (28). Both, metagenomics and 16S rRNA gene
525 sequencing, indicated that *Nitrosomonas* and *Nitrospira*-like bacteria were the only known
526 nitrifiers present in the systems investigated. This is consistent with numerous studies
527 (26,35,89,90) suggesting the specific adaptation of these genera to the wastewater environment.
528 The majority of the *Nitrosomonas*-like AOB MAGs, within and between systems, had AAI values
529 below 85% (Figure 4) indicating that this functional group exhibited high species level diversity.
530 Meanwhile, *Nitrospira*-like MAGs within the same lineage reported AAI values above 85% in
531 most cases. The high population level diversity of the *Nitrosomonas*-like bacteria was also
532 confirmed with full length 16S rRNA gene sequencing (Figure S3B) and corroborates earlier
533 studies on their diversity in wastewater (26,90,91). Although *N. europaea*, *N. oligotropha* and the
534 *Nitrosomonas*-like clusters closely associated with *N. communis/nitrosa* are frequently the most
535 abundant groups in wastewater, different *Nitrosomonas* populations dominate different processes
536 and the extent of their diversity is system dependent (26). For example, *N. europaea* are often the
537 major AOB in high ammonia concentration environments (e.g., anammox reactors) due to low
538 ammonia affinity and low DO level adaptation (91). The observed dominance of *N. oligotropha*
539 and *N. communis* in this study are consistent with previous studies of WWTPs with relatively
540 lower ammonia concentrations (26,91,92).

541
542 Comammox bacteria exhibited significantly lower population level diversity as compared to strict
543 AOB and NOB (Figure 4). All comammox *Nitrospira* MAGs, independent of the system they were
544 recovered from, belong to *Nitrospira* lineage II and were closely associated with *Ca. Nitrospira*
545 *nitrosa*. These findings are consistent with several other studies that have found *Ca. Nitrospira*
546 *nitrosa*-like bacteria as the unique or principal comammox bacteria present in their laboratory and
547 full-scale wastewater treatment systems (11,15,17,20,22,75,76), both using metagenomics and
548 gene targeted analyses. In contrast, other engineered and natural habitats typically harbor multiple
549 co-existing species across the two primary clades (i.e., comammox clade A and B). For example,
550 a study of multiple groundwater-fed sand filters recovered 12 comammox MAGs (5 clade A and
551 7 clade B) and revealed the co-occurrence of at least five comammox species (10 on average) per
552 system (34). Other studies in soils, sediments, lakes, drinking water, groundwater, and estuaries
553 have also reported the co-existence of multiple comammox species (9,73,93,94). However, this
554 co-existence of comammox populations across clades does not seem to be prevalent in wastewater
555 treatment systems. For instance, a study of 8 WWTPs found 9 of 14 comammox OTUs were
556 associated with the *Ca. Nitrospira nitrosa* cluster and accounted for 94.34% of all comammox
557 *amoA* sequences (17) with different OTUs representing the dominant comammox species in
558 different systems. Moreover, Beach and Noguera (75) reported the dominance of *Ca. Nitrospira*
559 *nitrosa* in five low DO nitrogen removal bioreactors using qPCR with species-specific primers.
560 These findings suggest that the diversity of comammox bacteria in WWTPs is lower than in other
561 habitats where several comammox species co-exist.

562

563 In contrast, Spasov et al. (16) reported the detection of multiple comammox MAGs in samples
564 collected from rotating biological contactors (RBCs) used for tertiary treatment at a municipal
565 WWTP in Ontario, Canada. The difference in population diversity between that study and other
566 WWTP investigations, including this one, may be associated with ammonia availability.
567 Specifically, tertiary treatment systems are typically employed as a polishing step; they are
568 designed and operated to remove low residual amounts of nitrogen. The influent ammonium
569 concentrations reported for the RBC tertiary treatment system range from 0.2 to 16.3 μM (16),
570 while the three systems in this study have influent ammonia concentrations between 2 and 3 mM.
571 The comammox *Nitrospira* MAGs with overall higher abundances in the tertiary plant (RBC001
572 and RBC083) were less abundant at the beginning of the train (higher ammonium concentrations)
573 than at the end of the train (lower ammonium concentration). In contrast, the abundance of
574 RBC035, the only MAG phylogenetically associated with *Ca. Nitrospira nitrosa* (ANI >95%),
575 decreased in abundance with decreasing ammonium concentrations. The high comammox
576 diversity in the RBCs and other ammonium-limited systems suggest that ammonium concentration
577 and/or flux through the system may be a key factor driving comammox bacteria diversity in
578 nitrification systems. In fact, Palomo et al. (34) reported that the influent ammonium concentration
579 was the key explanatory variable associated with comammox bacterial diversity in 12
580 groundwater-fed sand filters when higher comammox population diversity was detected in systems
581 receiving lower ammonium concentrations. *Ca. Nitrospira inopinata* and *Ca. Nitrospira kreftii*,
582 two comammox species with known kinetic parameters, have high apparent ammonia affinities
583 (95,96); assuming this is a conserved trait in comammox bacteria, this explains their prevalence
584 and diversity in ammonium-limited systems. However, the kinetic parameters (i.e., affinity
585 constant, growth rate) of other comammox species especially of the *Ca. Nitrospira nitrosa* clade

586 have not yet been reported, and our findings suggest that at least this comammox clade may have
587 substantially different kinetic traits that allow it to thrive in systems with higher ammonium
588 concentrations. Low DO has also been associated with the prevalence of comammox bacteria over
589 canonical nitrifiers in wastewater systems (15,75). However, the high DO concentrations (>2
590 mg/L) in this and other studies (17,20) suggest DO is not a crucial factor for the proliferation of
591 *Ca. Nitrospira nitrosa* in nitrogen removal systems.

592

593 **4.3 Comammox bacteria not only exhibit low diversity at the species/population level but** 594 **also at the strain/intra-population level across wastewater systems**

595 The high abundance and microdiversity of *Nitrospira*-NOB have been previously reported in
596 wastewater and other engineered and natural environments (4,33,35). However, it is unclear
597 whether the high microdiversity of *Nitrospira* is associated with high functional diversity or is
598 indicative of the coexistence of functionally identical *Nitrospira* with allelic diversity. Gruber-
599 Dorninger et al. (35) reported variable responses of closely related bacteria within *Nitrospira*
600 lineage I (sequence identities ranging from 95.8 to 99.6%) to different nitrite availabilities.
601 Similarly, ecological niche partitioning was also identified as potential mechanisms for co-
602 existence of three *Nitrospira* lineage I strains that used formate under different conditions: two
603 used formate when incubated with nitrite and ammonia, respectively, while the third used formate
604 efficiently as the sole substrate (35). Although these results shed light on the possible niche
605 partitioning of *Nitrospira* sub-species, this does not shed light on whether the microdiversity in
606 our study represents coexisting *Nitrospira* strains with slight functional differences. Although
607 functional implications of such high microdiversity are not clear, the persistence of high intra-
608 population diversity may suggest it plays an important role in the distribution and success of

609 *Nitrospira* populations in wastewater systems. A few studies in other environments have shown
610 that persistent populations exhibit increased intra-population sequence diversity (7,32). A study in
611 a saltern pond suggested that the ecologically important genes of the major archaeal sequence-
612 discrete population were carried by distinct sub-populations (strains), indicating that the adaptation
613 to different salinity concentrations had led to sub-population differentiation and speciation (niche
614 partitioning) (82). Moreover, a study in eight different temperate bog lakes concluded that high
615 microdiversity is associated with the maintenance of functional microbial communities during
616 changes in environmental conditions (36).

617

618 Contrastingly, comammox bacteria not only exhibited an unexpected low population-level
619 diversity within and between systems but also showed significantly lower microdiversity as
620 compared to strict NOB within the genus *Nitrospira*. These results may indicate specific
621 adaptations of comammox bacteria within the *Ca. Nitrospira nitrosa* cluster to the wastewater
622 environment and particularly secondary treatment systems. Palomo et al. (34) observed a negative
623 correlation between the species-level diversity of comammox *Nitrospira* and ammonium
624 concentrations. However, that was not the case for microdiversity, suggesting that different
625 mechanisms may shape inter- versus intra-population diversities, or that the range of ammonium
626 concentrations in the investigated drinking water systems was too narrow to capture any
627 underlying associations. For instance, the comammox nucleotide diversity in our study ranged
628 from 0.002 to 0.008, while this was from ~0.005 to ~0.013 for the MAGs obtained from these
629 drinking water systems, with only three of the twelve comammox MAGs presenting nucleotide
630 diversities lower than 0.008 (34). This may suggest that comammox microdiversity is also
631 associated with ammonium concentrations and/or flux through the system, with the observed low

632 microdiversity in wastewater systems associated with the higher prevailing ammonium
633 concentrations.

634 Despite the broad detection of comammox bacteria in WWTPs, their role and process relevance
635 are as yet unclear. In contrast to other environments, comammox *Nitrospira* present very low
636 population and intra-population diversity levels in wastewater treatment systems for nitrogen
637 removal. This observed lack of diversity and, consequently, lack of functional redundancy may
638 influence the feasibility of potential design and operational strategies relying primarily on
639 comammox *Nitrospira* for nitrogen removal. Although ammonium concentration and/or
640 availability apparently influences the diversity of comammox bacteria, further studies are
641 necessary to determine other factors driving the success of their clonal community in wastewater.
642 Moreover, studies are required to estimate the activity of comammox bacteria, specifically *Ca.*
643 *Nitrospira nitrosa*, and to assess their contribution to nitrification in full-scale nitrogen removal
644 systems. It is also important to note that the intra-population diversity cannot be solely studied
645 with shotgun metagenomic methods since high microdiversity can prevent robust assembly of
646 individual genomes (82). Therefore, more accurate techniques are needed to obtain high-quality
647 strain-level genome assemblies for *Nitrospira* and other highly microdiverse bacteria.

648

649 **4 Acknowledgements**

650 This research was supported by the National Science Foundation (Award number: 1703089) and
651 the Netherlands Organization for Scientific Research (016.Vidi.189.050). The authors also
652 acknowledge the operational personnel for assistance with sampling and process data sharing. The
653 authors also thanks A.M. Eren for assistance with metagenomic co-assemblies.

654

655 **5 Data availability**

656 All raw sequencing data from Illumina and Nanopore platforms and nitrifiers MAGs are

657 available on NCBI under bioproject number PRJNA846349.

658 6 References

- 659 1. Klotz MG, Stein LY. Nitrifier genomics and evolution of the nitrogen cycle. *FEMS*
660 *Microbiol Lett.* 2008;278(2):146–56.
- 661 2. Daims H, Lebedeva E v, Pjevac P, Han P, Herbold C, Albertsen M, et al. Complete
662 nitrification by *Nitrospira* bacteria. *Nature.* 2015;528(7583):504–9.
- 663 3. Pinto AJ, Marcus DN, Ijaz UZ, Bautista-de Lose Santos QM, Dick GJ, Raskin L.
664 Metagenomic evidence for the presence of comammox *Nitrospira*-like bacteria in a
665 drinking water system. *Mosphere.* 2016;1(1): e00054–15.
- 666 4. Palomo A, Jane Fowler S, Gülay A, Rasmussen S, Sicheritz-Ponten T, Smets BF.
667 Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel
668 physiology of *Nitrospira* spp. *ISME J.* 2016;10(11):2569–81.
- 669 5. van Kessel MAHJ, Speth DR, Albertsen M, Nielsen PH, den Camp HJM, Kartal B, et al.
670 Complete nitrification by a single microorganism. *Nature.* 2015;528(7583):555–9.
- 671 6. Hu HW, He JZ. Comammox—a newly discovered nitrification process in the terrestrial
672 nitrogen cycle. *Journal of Soils and Sediments.* 2017;17(12):2709–17.
- 673 7. Orellana LH, ben Francis T, Krüger K, Teeling H, Müller MC, Fuchs BM, et al. Niche
674 differentiation among annually recurrent coastal Marine Group II Euryarchaeota. *ISME J.*
675 2019;13(12):3024–36.
- 676 8. Wang Y, Ma L, Mao Y, Jiang X, Xia Y, Yu K, et al. Comammox in drinking water
677 systems. *Water Res.* 2017; 116:332–41.
- 678 9. Pjevac P, Schauburger C, Poghosyan L, Herbold CW, van Kessel MAHJ, Daebeler A, et
679 al. *AmoA*-targeted polymerase chain reaction primers for the specific detection and
680 quantification of comammox *Nitrospira* in the environment. *Front Microbiol.* 2017;
681 8:1508.
- 682 10. Annavajhala MK, Kapoor V, Santo-Domingo J, Chandran K. Comammox functionality
683 identified in diverse engineered biological wastewater treatment systems. *Environ Sci*
684 *Technol Lett.* 2018;5(2):110–6.
- 685 11. Camejo PY, Santo Domingo J, McMahon KD, Noguera DR. Genome-enabled insights
686 into the ecophysiology of the comammox bacterium “*Candidatus Nitrospira nitrosa*.”
687 *Msystems.* 2017;2(5): e00059–17.
- 688 12. Chao Y, Mao Y, Yu K, Zhang T. Novel nitrifiers and comammox in a full-scale hybrid
689 biofilm and activated sludge reactor revealed by metagenomic approach. *Appl Microbiol*
690 *Biotechnol.* 2016;100(18):8225–37.
- 691 13. Fan XY, Gao JF, Pan KL, Li DC, Dai HH. Temporal dynamics of bacterial communities
692 and predicted nitrogen metabolism genes in a full-scale wastewater treatment plant. *Rsc*
693 *Advances.* 2017;7(89):56317–27.
- 694 14. Gonzalez-Martinez A, Rodriguez-Sanchez A, van Loosdrecht MC, Gonzalez-Lopez J,
695 Vahala R. Detection of comammox bacteria in full-scale wastewater treatment bioreactors
696 using tag-454-pyrosequencing. *Environmental Science and Pollution Research.*
697 2016;23(24):25501–11.
- 698 15. Roots P, Wang Y, Rosenthal AF, Griffin JS, Sabba F, Petrovich M, et al. Comammox
699 *Nitrospira* are the dominant ammonia oxidizers in a mainstream low dissolved oxygen
700 nitrification reactor. *Water Res.* 2019; 157:396–405.
- 701 16. Spasov E, Tsuji JM, Hug LA, Doxey AC, Sauder LA, Parker WJ, et al. High functional
702 diversity among *Nitrospira* populations that dominate rotating biological contactor

- 703 microbial communities in a municipal wastewater treatment plant. *ISME J.*
704 2020;14(7):1857–72.
- 705 17. Wang M, Huang G, Zhao Z, Dang C, Liu W, Zheng M. Newly designed primer pair
706 revealed dominant and diverse comammox amoA gene in full-scale wastewater treatment
707 plants. *Bioresour Technol.* 2018; 270:580–7.
- 708 18. Xia F, Wang JG, Zhu T, Zou B, Rhee SK, Quan ZX. Ubiquity and diversity of complete
709 ammonia oxidizers (comammox). *Appl Environ Microbiol.* 2018;84(24): e01390–18.
- 710 19. Yang Y, Daims H, Liu Y, Herbold CW, Pjevac P, Lin JG, et al. Activity and metabolic
711 versatility of complete ammonia oxidizers in full-scale wastewater treatment systems.
712 *Mbio.* 2020;11(2): e03175–19.
- 713 20. Xu S, Chai W, Xiao R, Smets BF, Palomo A, Lu H. Survival strategy of comammox
714 bacteria in a wastewater nutrient removal system with sludge fermentation liquid as
715 additional carbon source. *Science of The Total Environment.* 2022; 802:149862.
- 716 21. Palomo A, Dechesne A, Smets BF. Genomic profiling of *Nitrospira* species reveals
717 ecological success of comammox *Nitrospira*. *BioRxiv.* 2019;612226.
- 718 22. Zhao Z, Huang G, He S, Zhou N, Wang M, Dang C, et al. Abundance and community
719 composition of comammox bacteria in different ecosystems by a universal primer set.
720 *Science of The Total Environment.* 2019; 691:146–55.
- 721 23. Xia F, Wang JG, Zhu T, Zou B, Rhee SK, Quan ZX. Ubiquity and diversity of complete
722 ammonia oxidizers (comammox). *Appl Environ Microbiol.* 2018;84(24): e01390–18.
- 723 24. Gottshall EY, Bryson SJ, Cogert KI, Landreau M, Sedlacek CJ, Stahl DA, et al. Sustained
724 nitrogen loss in a symbiotic association of Comammox *Nitrospira* and Anammox bacteria.
725 *Water Research.* 2021; 202:117426.
- 726 25. Li D, Fang F, Liu G. Efficient nitrification and low-level N₂O emission in a weakly acidic
727 bioreactor at low dissolved-oxygen levels are due to comammox. *Appl Environ Microbiol.*
728 2021;87(11): e00154–21.
- 729 26. Gao JF, Luo X, Wu GX, Li T, Peng YZ. Quantitative analyses of the composition and
730 abundance of ammonia-oxidizing archaea and ammonia-oxidizing bacteria in eight full-
731 scale biological wastewater treatment plants. *Bioresour Technol.* 2013; 138:285–96.
- 732 27. Ofileru ID, Lunn M, Curtis TP, Wells GF, Criddle CS, Francis CA, et al. Combined
733 niche and neutral effects in a microbial wastewater treatment community. *Proceedings of*
734 *the National Academy of Sciences.* 2010;107(35):15345–50.
- 735 28. Siripong S, Rittmann BE. Diversity study of nitrifying bacteria in full-scale municipal
736 wastewater treatment plants. *Water Res.* 2007;41(5):1110–20.
- 737 29. Konopka A, Lindemann S, Fredrickson J. Dynamics in microbial communities: unraveling
738 mechanisms to identify principles. *ISME J.* 2015;9(7):1488–95.
- 739 30. Schloter M, Leubhn M, Heulin T, Hartmann A. Ecology and evolution of bacterial
740 microdiversity. *FEMS Microbiol Rev.* 2000;24(5):647–60.
- 741 31. Cotto I, Dai Z, Huo L, Anderson CL, Vilardi KJ, Ijaz U, et al. Long solids retention times
742 and attached growth phase favor prevalence of comammox bacteria in nitrogen removal
743 systems. *Water Res.* 2020; 169:115268.
- 744 32. Meziti A, Tsementzi D, Rodriguez-R LM, Hatt JK, Karayanni H, Kormas KA, et al.
745 Quantifying the changes in genetic diversity within sequence-discrete bacterial
746 populations across a spatial and temporal riverine gradient. *ISME J.* 2019;13(3):767–79.

- 747 33. Gülay A, Musovic S, Albrechtsen HJ, Al-Soud WA, Sørensen SJ, Smets BF. Ecological
748 patterns, diversity and core taxa of microbial communities in groundwater-fed rapid
749 gravity filters. *ISME J.* 2016;10(9):2209–22.
- 750 34. Palomo A, Dechesne A, Cordero OX, Smets BF. Evolutionary ecology of natural
751 comammox *Nitrospira* populations. *Msystems.* 2022;7(1): e01139–21.
- 752 35. Gruber-Dorninger C, Pester M, Kitzinger K, Savio DF, Loy A, Rattei T, et al.
753 Functionally relevant diversity of closely related *Nitrospira* in activated sludge. *ISME J.*
754 2015;9(3):643–55.
- 755 36. García-García N, Tamames J, Linz AM, Pedrós-Alió C, Puente-Sánchez F.
756 Microdiversity ensures the maintenance of functional microbial communities under
757 changing environmental conditions. *ISME J.* 2019;13(12):2969–83.
- 758 37. Bertrand D, Shaw J, Kalathiyappan M, Ng AHQ, Kumar MS, Li C, et al. Hybrid
759 metagenomic assembly enables high-resolution analysis of resistance determinants and
760 mobile elements in human microbiomes. *Nat Biotechnol.* 2019;37(8):937–44.
- 761 38. Mantere T, Kersten S, Hoischen A. Long-read sequencing emerging in medical genetics.
762 *Front Genet.* 2019; 10:426.
- 763 39. Singleton CM, Petriglieri F, Kristensen JM, Kirkegaard RH, Michaelsen TY, Andersen
764 MH, et al. Connecting structure to function with the recovery of over 1000 high-quality
765 metagenome-assembled genomes from activated sludge using long-read sequencing. *Nat*
766 *Commun.* 2021;12(1):1–13.
- 767 40. Frank JA, Reich CI, Sharma S, Weisbaum JS, Wilson BA, Olsen GJ. Critical evaluation of
768 two primers commonly used for amplification of bacterial 16S rRNA genes. *Applied and*
769 *Environmental Microbiology.* 2008 Apr;74(8):2461–70.
- 770 41. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
771 High-resolution sample inference from Illumina amplicon data. *Nat Methods.*
772 2016;13(7):581–3.
- 773 42. Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, et al. High-throughput
774 amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution.
775 *Nucleic Acids Res.* 2019;47(18): e103–e103.
- 776 43. Wright ES. RNAconTest: comparing tools for noncoding RNA multiple sequence
777 alignment based on structural consistency. *RNA.* 2020;26(5):531–40.
- 778 44. Stackebrandt E. Taxonomic parameters revisited: tarnished gold standards. *Microbiol*
779 *Today.* 2006; 33:152–5.
- 780 45. Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer KH, et al. Uniting the
781 classification of cultured and uncultured bacteria and archaea using 16S rRNA gene
782 sequences. *Nature Reviews Microbiology.* 2014;12(9):635–45.
- 783 46. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
784 throughput. *Nucleic Acids Res.* 2004;32(5):1792–7.
- 785 47. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
786 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.*
787 2015;32(1):268–74.
- 788 48. Price MN, Dehal PS, Arkin AP. FastTree 2—approximately maximum-likelihood trees for
789 large alignments. *PLoS One.* 2010;5(3): e9490.
- 790 49. Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial
791 communities. *Appl Environ Microbiol.* 2005;71(12):8228–35.
- 792 50. Wickham H. *ggplot2: elegant graphics for data analysis.* springer; 2016.

- 793 51. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
794 *Bioinformatics*. 2018;34(17): i884–i890.
- 795 52. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
796 SAMtools and BCFtools. *Gigascience*. 2021;10(2): giab008.
- 797 53. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
798 features. *Bioinformatics*. 2010;26(6):841–2.
- 799 54. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile
800 metagenomic assembler. *Genome Res*. 2017;27(5):824–34.
- 801 55. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an
802 advanced analysis and visualization platform for 'omics data. *PeerJ*. 2015;3: e1319.
- 803 56. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.
804 *bioinformatics*. 2009;25(14):1754–60.
- 805 57. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for
806 genome assemblies. *Bioinformatics*. 2013;29(8):1072–5.
- 807 58. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive
808 binning algorithm for robust and efficient genome reconstruction from metagenome
809 assemblies. *PeerJ*. 2019;7: e7359.
- 810 59. Alneberg J, Bjarnason BS, Bruijn I de, Schirmer M, Quick J, Ijaz UZ, et al. Binning
811 metagenomic contigs by coverage and composition. *Nature Methods*. 2014
812 Apr;11(11):1144–6.
- 813 60. Wu YW, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to
814 recover genomes from multiple metagenomic datasets. *Bioinformatics*. 2016;32(4):605–7.
- 815 61. Warren RL, Coombe L, Mohamadi H, Zhang J, Jaquish B, Isabel N, et al. ntEdit: scalable
816 genome sequence polishing. *Bioinformatics*. 2019;35(21):4430–2.
- 817 62. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
818 quality of microbial genomes recovered from isolates, single cells, and metagenomes.
819 *Genome Res*. 2015;25(7):1043–55.
- 820 63. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarszewski A, Chaumeil PA, et al. A
821 standardized bacterial taxonomy based on genome phylogeny substantially revises the tree
822 of life. *Nat Biotechnol*. 2018;36(10):996–1004.
- 823 64. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
824 prokaryotic gene recognition and translation initiation site identification. *BMC*
825 *Bioinformatics*. 2010;11(1):1–11.
- 826 65. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference
827 resource for gene and protein annotation. *Nucleic Acids Res*. 2016;44(D1): D457–D462.
- 828 66. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.
829 KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score
830 threshold. *Bioinformatics*. 2020;36(7):2251–2.
- 831 67. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of
832 genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat*
833 *Microbiol*. 2018;3(7):836–43.
- 834 68. Woodcroft B, Newell R. CoverM (Version 0.6.1). 2021.
- 835 69. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput
836 ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun*.
837 2018;9(1):1–8.

- 838 70. Parks D, Skennerton C, Brown C. CompareM. <https://github.com/dparks1134/CompareM>;
839 2020.
- 840 71. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF.
841 inStrain profiles population microdiversity from metagenomic data and sensitively detects
842 shared microbial strains. *Nature Biotechnology*. 2021;39(6):727–36.
- 843 72. Uritskiy G v, DiRuggiero J, Taylor J. MetaWRAP—a flexible pipeline for genome-
844 resolved metagenomic data analysis. *Microbiome*. 2018;6(1):1–13.
- 845 73. Xu Y, Lu J, Wang Y, Liu G, Wan X, Hua Y, et al. Diversity and abundance of
846 comammox bacteria in the sediments of an urban lake. *Journal of Applied Microbiology*.
847 2020;128(6):1647–57.
- 848 74. Shi Y, Jiang Y, Wang S, Wang X, Zhu G. Biogeographic distribution of comammox
849 bacteria in diverse terrestrial habitats. *Science of the Total Environment*. 2020;
850 717:137257.
- 851 75. Beach NK, Noguera DR. Design and assessment of species-level qPCR primers targeting
852 comammox. *Front Microbiol*. 2019; 10:36.
- 853 76. Zheng M, Wang M, Zhao Z, Zhou N, He S, Liu S, et al. Transcriptional activity and
854 diversity of comammox bacteria as a previously overlooked ammonia oxidizing
855 prokaryote in full-scale wastewater treatment plants. *Science of the Total Environment*.
856 2019; 656:717–22.
- 857 77. Zhou X, Li B, Wei J, Ye Y, Xu J, Chen L, et al. Temperature Influenced the Comammox
858 Community Composition in Drinking Water and Wastewater Treatment Plants. *Microbial
859 Ecology*. 2021;82(4):870–84.
- 860 78. Koops HP, Purkhold U, Pommerening-Röser A, Timmermann G, Wagner M. The
861 lithoautotrophic ammonia-oxidizing bacteria. *Bergey’s manual of systematic bacteriology*,
862 volume two: the proteobacteria, part A introductory essays, Springer. 2005;141–7.
- 863 79. Pommerening-Röser A, Rath G, Koops HP. Phylogenetic diversity within the genus
864 *Nitrosomonas*. *Systematic and Applied Microbiology*. 1996;19(3):344–51.
- 865 80. Koops HP, Stehr G, others. Classification of eight new species of ammonia-oxidizing
866 bacteria: *Nitrosomonas communis* sp. nov., *Nitrosomonas ureae* sp. nov., *Nitrosomonas
867 aestuarii* sp. nov., *Nitrosomonas marina* sp. nov., *Nitrosomonas nitrosa* sp. nov.,
868 *Nitrosomonas eutropha* sp. nov., *Nitrosomonas oligotropha* sp. nov. and *Nitrosomonas
869 halophila* sp. nov. *Microbiology (N Y)*. 1991;137(7):1689–99.
- 870 81. Navada S, Knutsen MF, Bakke I, Vadstein O. Nitrifying biofilms deprived of organic
871 carbon show higher functional resilience to increases in carbon supply. *Sci Rep*.
872 2020;10(1):1–11.
- 873 82. Konstantinidis KT, Viver T, Conrad RE, Venter SN, Rossello-Mora R. Solar salterns as
874 model systems to study the units of bacterial diversity that matter for ecosystem
875 functioning. *Curr Opin Biotechnol*. 2022; 73:151–7.
- 876 83. Luo C, Rodriguez-r LM, Konstantinidis KT. MyTaxa: an advanced taxonomic classifier
877 for genomic and metagenomic sequences. *Nucleic Acids Res*. 2014;42(8): e73–e73.
- 878 84. Sczyrba A, Hofmann P, Belmann P, Koslicki D, Janssen S, Dröge J, et al. Critical
879 assessment of metagenome interpretation—a benchmark of metagenomics software. *Nat
880 Methods*. 2017;14(11):1063–71.
- 881 85. de Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J, et al. Comparison
882 of long-read sequencing technologies in the hybrid assembly of complex bacterial
883 genomes. *Microb Genom*. 2019;5(9).

- 884 86. Xie H, Yang C, Sun Y, Igarashi Y, Jin T, Luo F. PacBio long reads improve metagenomic
885 assemblies, gene catalogs, and genome binning. *Front Genet.* 2020;1077.
- 886 87. Brown CL, Keenum IM, Dai D, Zhang L, Vikesland PJ, Pruden A. Critical evaluation of
887 short, long, and hybrid assembly for contextual analysis of antibiotic resistance genes in
888 complex environmental metagenomes. *Sci Rep.* 2021;11(1):1–12.
- 889 88. Dilhari A, Sampath A, Gunasekara C, Fernando N, Weerasekara D, Sissons C, et al.
890 Evaluation of the impact of six different DNA extraction methods for the representation of
891 the microbial community associated with human chronic wound infections using a gel-
892 based DNA profiling method. *Amb Express.* 2017;7(1):1–11.
- 893 89. Dionisi HM, Layton AC, Harms G, Gregory IR, Robinson KG, Sayler GS. Quantification
894 of *Nitrosomonas oligotropha*-like ammonia-oxidizing bacteria and *Nitrospira* spp. from
895 full-scale wastewater treatment plants by competitive PCR. *Appl Environ Microbiol.*
896 2002;68(1):245–53.
- 897 90. Zhang T, Ye L, Tong AHY, Shao MF, Lok S. Ammonia-oxidizing archaea and ammonia-
898 oxidizing bacteria in six full-scale wastewater treatment bioreactors. *Appl Microbiol*
899 *Biotechnol.* 2011;91(4):1215–25.
- 900 91. Gao J, Luo X, Wu G, Li T, Peng Y. Abundance and diversity based on *amoA* genes of
901 ammonia-oxidizing archaea and bacteria in ten wastewater treatment systems. *Appl*
902 *Microbiol Biotechnol.* 2014;98(7):3339–54.
- 903 92. Wells GF, Park HD, Yeung CH, Eggleston B, Francis CA, Criddle CS. Ammonia-
904 oxidizing communities in a highly aerated full-scale activated sludge bioreactor:
905 betaproteobacterial dynamics and low relative abundance of Crenarchaea. *Environ*
906 *Microbiol.* 2009;11(9):2310–28.
- 907 93. Vilaridi KJ, Cotto I, Rivera MS, Dai Z, Anderson CL, Pinto A. Comammox *Nitrospira*
908 bacteria outnumber canonical nitrifiers irrespective of electron donor mode and
909 availability. *FEMS Microbiol Ecol.* 2022;98(4): fiac032.
- 910 94. Wang X, Lu L, Zhou X, Tang X, Kuang L, Chen J, et al. Niche differentiation of
911 comammox *Nitrospira* in the mudflat and reclaimed agricultural soils along the north
912 branch of Yangtze River estuary. *Front Microbiol.* 2021;3482.
- 913 95. Sakoula D, Koch H, Frank J, Jetten MSM, van Kessel MAHJ, Lückner S. Enrichment and
914 physiological characterization of a novel comammox *Nitrospira* indicates ammonium
915 inhibition of complete nitrification. *ISME Journal.* 2021 Apr 1;15(4):1010–24.
- 916 96. Kits KD, Sedlacek CJ, Lebedeva E v, Han P, Bulaev A, Pjevac P, et al. Kinetic analysis of
917 a complete nitrifier reveals an oligotrophic lifestyle. *Nature.* 2017;549(7671):269–72.
- 918