

1

2

3

4 Expression of the Eight GABA_A Receptor α Subunits in the
5 Developing Zebrafish Central Nervous System

6

7

8 Bryan Monesson-Olson^{1,2}, Jon J. McClain², Abigail E. Case², Hanna E. Dorman², Daniel R.
9 Turkewitz³, Aaron B. Steiner³ and Gerald B. Downes^{1, 2*}

10

11

12

13 ¹Neuroscience and Behavior Graduate Program

14 ²Biology Department, University of Massachusetts, Amherst, MA 01003, USA

15 ³Department of Biology and Health Sciences, Pace University, Pleasantville, NY 10570, USA

16

17

18 *Corresponding author:

19 Gerald B. Downes

20 Email: gbdownes@bio.umass.edu

21 **ABSTRACT**

22 GABA is a robust regulator of both developing and mature neural networks. It exerts
23 many of its effects through GABA_A receptors, which are heteropentamers assembled from a
24 large array of subunits encoded by distinct genes. In mammals, there are 19 different GABA_A
25 subunit types, which are divided into the α , β , γ , δ , ϵ , π , θ and ρ subfamilies. The immense
26 diversity of GABA_A receptors is not fully understood. However, it is known that specific isoforms,
27 with their distinct biophysical properties and expression profiles, tune responses to GABA.
28 Although larval zebrafish are well-established as a model system for neural circuit analysis, little
29 is known about GABA_A receptors diversity and expression in this system. Here, using database
30 analysis, we show that the zebrafish genome contains at least 23 subunits. All but the
31 mammalian θ and ϵ subunits have at least one zebrafish ortholog, while five mammalian GABA_A
32 subunits have two zebrafish orthologs. Zebrafish contain one subunit, β 4, which does not have
33 a clear mammalian ortholog. Similar to mammalian GABA_A receptors, the zebrafish α subfamily
34 is the largest and most diverse of the subfamilies. In zebrafish there are eight α subunits, and
35 RNA *in situ* hybridization across early zebrafish development revealed that they demonstrate
36 distinct patterns of expression in the brain, spinal cord, and retina. Some subunits were very
37 broadly distributed, whereas others were restricted to small populations of cells. Subunit-specific
38 expression patterns were similar between zebrafish, frogs and rodents, which suggests that the
39 roles of different GABA_A receptor isoforms are largely conserved among vertebrates. This study
40 provides a platform to examine isoform specific roles of GABA_A receptors within zebrafish
41 neural circuits and it highlights the potential of this system to better understand the remarkable
42 heterogeneity of GABA_A receptors.

43

44 INTRODUCTION

45 Neural networks throughout the central nervous system rely upon a diversity of
46 neurotransmitter systems for both their initial formation and mature function. GABA is the major
47 inhibitory neurotransmitter throughout most of the mature nervous system [1]. It exerts its effects
48 through its receptors, which are divided into two classes. GABA_A receptors are ligand-gated
49 chloride channels responsible for most of the rapid effects of GABA, while GABA_B receptors are
50 heterotrimeric G-protein coupled receptors. In mammalian systems, GABA_A receptors
51 demonstrate incredible subtype diversity and are targets for classes of clinically important drugs,
52 such as benzodiazepines and barbituates [2, 3]. GABA_A receptors are heteropentamers
53 composed of various combinations of 19 different subunits: α 1-6, β 1-3, γ 1-3, δ , ϵ , π , θ and ρ 1-3
54 (previously referred to as GABA_C receptors) [4, 5]. Each subunit is encoded by a discrete gene
55 that is spatially and developmentally regulated to generate distinct, but often overlapping,
56 expression patterns [6-9]. Alternative splicing and RNA editing of some subunits further
57 increases the number of subtypes available [10]. Although this extensive receptor heterogeneity
58 is not fully understood, some subunits confer distinct biophysical and pharmacological
59 properties, interact with specific cytoplasmic proteins, and localize to specific subcellular
60 domains [11-13]. Ultimately, this receptor diversity provides a capacity to tailor responses to
61 GABA within neural circuits.

62 The α subunits play a prominent role in GABA_A receptor function. They are thought to be
63 essential components of 'classic' GABA_A receptors, which exclude the pharmacologically
64 distinct receptors composed from ρ subunits [3]. Exogenous expression studies and analysis of
65 native receptors indicate that each receptor contains two α subunits which, along with β
66 subunits, form the GABA binding site. α subunits can also dictate isoform-selective
67 pharmacology. For example, receptors containing α 1, α 2, α 3 and α 5 subunits confer

68 benzodiazepine-sensitivity, whereas receptors that contain $\alpha 4$ and $\alpha 6$ subunits do not bind to
69 clinically used benzodiazepines [2]. In addition, α subunits regulate subcellular localization.
70 GABA receptors that contain $\alpha 1$, $\alpha 2$, and $\alpha 3$ demonstrate a synaptic localization and mediate
71 transient or phasic inhibition, while receptors that contain $\alpha 4$, $\alpha 5$ and $\alpha 6$ are principally
72 extrasynaptic and mediate tonic inhibition [14, 15]. Through both transient and tonic inhibition,
73 these receptors mediate the majority of inhibition in neural circuits throughout the vertebrate
74 brain.

75 Developing zebrafish have several features that make them well-suited to study the
76 formation and function of neural circuits [16, 17]. First, zebrafish embryos and larvae develop
77 external to the mother and are optically transparent. These features make the central nervous
78 system easily accessible throughout development and particularly amenable to optical
79 physiological approaches, such as optogenetics or calcium imaging. Second, the central
80 nervous system contains fewer cells compared to mammalian preparations, yet many cell types
81 and mechanisms are conserved among vertebrates. Lastly, larval zebrafish develop rapidly,
82 therefore many sensory and motor systems are present and functional within five days post-
83 fertilization. Combined, these features have helped establish developing zebrafish as a
84 prominent system to examine neural circuit formation and function.

85 GABA_A receptors are expressed in larval zebrafish and essential for normal nervous
86 system function. For example, bath application of pentylenetetrazole, a convulsant and
87 noncompetitive GABA_A receptor antagonist, induces hyperactive behavior that can serve as a
88 model of seizures [18]. Similarly, insecticides known to target GABA_A receptors generate
89 hyperactive behavior in larvae [19]. Pharmacological blockade of GABA_A receptors has also
90 been shown to regulate the activity of specific cells, including retinal bipolar cells [20], optic
91 tectum neurons [21], Mitral cells in the olfactory bulb [22], spinal cord CoPA interneurons and
92 motoneurons [23], and the Mauthner cells, a pair of well-studied reticulospinal neurons found in

93 the amphibian and teleost hindbrain [24]. In these studies, the GABA_A receptor isoforms were
94 not identified. However, patch-clamp recordings of the Mauthner cells showed three distinct
95 GABA_A kinetic profiles, which were proposed to be caused by different receptor isoforms [25].

96 There is limited information about the heterogeneity of GABA_A receptors in zebrafish.
97 The most extensive study to date identified 23 GABA_A receptor subunits and examined their
98 expression broadly in adult zebrafish brain via RT-qPCR [26]. Despite playing prominent roles in
99 developing animals, little is known about the expression of GABA_A receptor subunits in
100 zebrafish embryos and larvae. $\alpha 1$ has been shown to be expressed at 50 and 72 hours post-
101 fertilization (hpf), $\gamma 2$ at 50 hpf, and $\alpha 2a$ (then named $\alpha 2$) from ~14 – 96 hpf. [27-29].
102 Unpublished $\alpha 6a$ expression data has also been deposited in the ZFIN database [30]. The
103 expression patterns of other GABA_A receptor subunits has not been reported.

104 In this study we performed phylogenetic analysis, which indicates that zebrafish contain
105 at least 23 different GABA_A receptor subunits. Although we observed some differences between
106 the zebrafish and mammalian GABA_A receptor subunit gene families, zebrafish contain
107 orthologs for most of the GABA_A receptor subunits found in mammals. To examine where and
108 when these GABA_A receptors are expressed in the zebrafish nervous system, we performed
109 whole-mount RNA *in situ* hybridization. We present the embryonic and larval expression of the
110 eight identified α subunit-encoding genes: *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, *gabra5*,
111 *gabra6a* and *gabra6b*. These data show that each subunit has a distinct expression pattern,
112 broadly similar to the reported expression of amphibian and rodent orthologs. Combined, these
113 results suggest that GABA_A isoform-specific roles are conserved among vertebrates, and they
114 establish a foundation to use the zebrafish system to better understand how GABA_A receptor
115 heterogeneity mediates neural circuit formation and function.

116

117 MATERIALS AND METHODS

118 Phylogenetic analysis

119 Homologous gene queries were performed using the National Center for Biotechnology
120 Information tBLASTn search tool. Protein sequences for each of the 19 mouse GABA_A receptor
121 subunits were used to search the zebrafish nucleotide collection. Matching zebrafish sequences
122 were evaluated to determine if they were splice variants from the same gene or generated from
123 different genes. When splice variants were identified, only the longest variant was used for
124 subsequent analysis. 19 mouse and 23 zebrafish protein sequences were used to assemble a
125 phylogenetic tree. The genes and Genbank accession numbers for the mouse sequences are:
126 *Gabra1*- NP_034380, *Gabra2*- NP_032092, *Gabra3*- NP_032093, *Gabra4*- NP_034381,
127 *Gabra5*- NP_795916, *Gabra6*- NP_00109311, *Gabrb1*- NP_032095, *Gabrb2*- NP_032096,
128 *Gabrb3*- NP_032097, *Gabrd*- NP_032098, *Gabre*- NP_059065, *Gabrg1*- NP_034382, *Gabrg2*-
129 NP_032099, *Gabrg3*- NP_032100, *Gaprp*- NP_666129, *Gabrq*- NP_065234, *Gabrr1*-
130 NP_032101, *Gabrr2*- NP_032102, and *Gabrr3*- NP_001074659. The Genbank accession
131 numbers for the zebrafish sequences are: *gabra1*- NM_001077326, *gabra2a*- XM_009307207,
132 *gabra2b*- XM_017359049, *gabra3*- XM_009295708, *gabra4*- NM_001017822, *gabra5*-
133 XM_001339475, *gabra6a*- XM_005173112, *gabra6b*- XM_002667357, *gabrb1*- XM_002664133,
134 *gabrb2*- NM_001024387, *gabrb3*- XM_005166079, *gabrb4*- XM_005173874, *gabrd*-
135 XM_695007, *gabrg1*- XM009307208, *gabrg2*- NM_001256250, *gabrg3*- XM_009302568, *gaprp*-
136 XM_005173293, *gabrr1*- NM_001025553, *gabrr2a*- NM_001045376, *gabrr2b*- XM_009294512,
137 *gabrr3a*- NM_001128760, *gabrr3b*- XM_009297450, *gabrz*- XM_005156247. To construct the
138 tree, a ClustalW alignment of amino acid sequences was performed using the Geneious
139 software package [31]. An unrooted PHYML consensus tree was then generated using 100
140 bootstrap replicates [32].

141 **Fish maintenance and breeding**

142 Zebrafish were raised and maintained using established husbandry procedures.
143 Embryos were kept at 28.5 °C in E3 media and staged according to morphological criteria [33].
144 All experiments were performed using Tuebingen (Tu) or tub longfin (TLF) wild type embryos.
145 From 0-24 hpf, embryos were grown in 0.01% Methylene Blue in E3 medium. At 24 hpf the
146 solution was changed to 0.0045% Phenylthiourea (PTU) in E3 to inhibit pigment formation. This
147 solution was changed every 24 hours until the fish were sacrificed. All animal protocols were
148 approved by the University of Massachusetts Institutional Animal Care and Use Committees
149 (IACUC).

150 **Whole-mount in situ hybridization**

151 Antisense digoxigenin probes were generated against *gabra1*, *gabra2a*, *gabra2b*,
152 *gabra3*, *gabra4*, *gabra5*, *gabra6a* and *gabra6b* (S1 Table). *In situ* probe synthesis utilized the
153 digoxigenin RNA labeling kit with SP6 or T7 RNA polymerases (Roche Diagnostics, Mannheim,
154 Germany). Whole-mount, colorimetric *in situ* hybridization was performed using established
155 protocols [34] and examined using a compound microscope (Zeiss, Thornwood, NY) attached to
156 a digital camera (Zeiss, Thornwood, NY). Cross sections were generated by imbedding *in situ*
157 hybridization stained embryos in 1.5% agar, 5% sucrose. The blocks were submerged in a 30%
158 sucrose solution overnight then cut into 20mm thick sections using a cryostat (Leica, Buffalo
159 Grove, IL). Representative larvae at 24, 48, and 96 hpf were mounted in either 100% glycerol or
160 dehydrated through a methanol series, equilibrated in a 2:1 benzylbenzoate/benzylalcohol
161 solution and mounted in a 10:1 Canada balsam/methyl saliylate misture. Images were captured
162 using a Zeiss Discovery v12 stereomicroscope with a 1.5x objective or a Zeiss Axioskop
163 Microscope with a 10x objective and an AxioCam MRc camera. All images were processed
164 using Adobe Photoshop, in which multiple focal planes were merged to produce single

165 representative images. The identify of neuroanatomical structures was determined using
166 zebrafish brain atlases [35-37].

167 **Reverse-Transcriptase PCR**

168 RT-PCR was used to analyze whether the *gabra3* and *gabra3*-like sequence were
169 portions of the same transcript. Primers designed against *gabra3*-like (Primer 1 5'
170 GGACGGCGGATGATGAGAAA-3') and *gabra3* (Primer 2 5'-CACGACCGTCCTGACTA-3',
171 Primer 3 5'-GTGGAGTAGATGTGGTGGGC-3') were used to amplify cDNA from wild-type
172 zebrafish larvae. RNA was extracted from 48 hpf zebrafish larvae using the RNAeasy kit
173 (Qiagen, Venlo, Netherlands) and reverse transcribed using the Accuscript RT-PCR system
174 (Stratagene). The PCR products were sequenced to confirm that they are portions from the
175 same transcript.

176

177 **RESULTS**

178 **The zebrafish GABA_A receptor subunit gene family is similar** 179 **in size and diversity to the mammalian GABA_A receptor gene** 180 **family.**

181 19 GABA_A receptor subunit genes have been identified in humans, mice, and rats [4, 5].
182 To establish the number and organization of zebrafish GABA_A receptor subunits, we used
183 mouse GABA_A receptor subunit amino acid sequences to query zebrafish genome databases.
184 We identified 23 zebrafish GABA_A receptor subunits, each encoded by a distinct gene.
185 Frequently splice variants were observed. In these cases, the principle splice isoform, as
186 indicated by the databases, was selected for analysis. 12 of the 19 mouse GABA_A receptor
187 subunits were found to have a single ortholog in zebrafish (Fig 1). Amino acid identities between
188 mouse and zebrafish orthologs ranged from 53.1 to 86.3%. Five additional mouse subunits, α 2,

189 $\alpha 6$, $\rho 2$, $\rho 3$, and π , each exhibit similarity with two subunits in zebrafish: *gabra2a* and *gabra2b*,
190 *gabra6a* and *gabra6b*, *gabrr2a* and *gabrr2b*, *gabrr3a* and *gabrr3b*, *gabrp* and *gabrz*,
191 respectively. The relatively high percentage of amino acid identity of these zebrafish subunits
192 with each other suggests that the zebrafish paralogs are duplicated genes. Duplicated genes
193 are often observed in zebrafish, and they are thought to be due to a whole genome duplication
194 within the teleost lineage [38, 39]. It was reported previously that there is a *gabra3* and an $\alpha 3$ -
195 like gene. While the $\alpha 3$ -like gene has been localized to chromosome 21, the genomic location of
196 the $\alpha 3$ -like gene is not known. Our database analysis suggested that these sequences could be
197 non-overlapping portions of the same gene, therefore RT-PCR was performed using primers
198 targeting $\alpha 3$ -like and *gabra3* sequences (S1 Fig). Transcripts were amplified that spanned the
199 region between *gabra3* and $\alpha 3$ -like sequences, indicating that these are different portions of the
200 same gene and that there is only one $\alpha 3$ encoding gene, *gabra3*.

201

202 **Fig 1. Phylogenetic analysis shows that the zebrafish GABA_A subunit gene family is**
203 **similar in size, diversity, and organization to the mouse GABA_A subunit gene family.**

204 Amino acid sequence alignments were used to generate a consensus tree using 100 bootstrap
205 replicates. The genes that encode the GABA_A subunits are shown at the tip of each branch and
206 bootstrap proportions are shown at the branch points. The zebrafish and mouse GABA_A subunit
207 sequences showed high amino acid identity and grouped into the α , β , γ , δ , π , and ρ
208 subfamilies. Most mouse GABA_A subunits have a single zebrafish ortholog, while six mouse
209 GABA_A subunits have two zebrafish orthologs, likely due to gene duplication.

210

211 There are two mouse GABA_A receptor subunits for which zebrafish orthologous are less
212 clear or have yet to be identified, the θ and ϵ subunits. The θ subunit, encoded by the gene
213 *Gabrq*, exhibits only 31.3% amino acid identity with zebrafish $\beta 4$, encoded by *gabrb4*, which is

214 the greatest percentage identity of this subunit with any zebrafish sequence. There are three β
215 subunits in mice compared to the four β subunits in zebrafish so it is possible that, despite the
216 low amino acid sequence identity, zebrafish $\beta 4$ is orthologous to the mouse θ subunit. No clear
217 zebrafish ortholog has yet to be identified for the mouse ϵ subunit, which is encoded by the
218 *Gabre* gene.

219

220 **α subunits demonstrate distinct expression patterns across** 221 **early zebrafish development**

222 Determining the temporal and spatial expression of GABA_A subunits is instrumental to
223 identify isoform-specific roles in zebrafish neural circuit development and function. Therefore,
224 given the large size of the α subunit subfamily and their prominent role in GABA_A receptor
225 function, we examined the expression of the eight α subunit encoding genes: *gabra1*, *gabra2a*,
226 *gabra2b*, *gabra3*, *gabra4*, *gabra6a* and *gabra6b*. Using whole-mount RNA *in situ* hybridization,
227 we determined their expression patterns at 24, 48, and 96 hpf. These time points span much of
228 larval zebrafish locomotor network development [40].

229 Expression of *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4* and *gabra5* was detected at 24
230 hpf (Fig 2). Neither *gabra6a* or *gabra6b* were detected at this developmental stage. *gabra2a*
231 and *gabra2b* were detected broadly and did not appear to be spatially restricted (Fig 2C-F). To
232 distinguish whether these gene transcripts were widespread or background due to the probe, a
233 second probe was generated for each gene (S1 Table). These probes yielded widespread
234 staining, very similar to the initial probes used, which suggests that *gabra2a* and *gabra2b* are
235 widely distributed. Consistent with these findings, a previous study reported that *gabra2a*
236 exhibits a diffuse, broad pattern of expression in larval zebrafish [28].

237

238 **Fig 2. Expression of GABA_A α subunits at 24 hpf.** *gabra1* (A, B), *gabra2a* (C, D), *gabra2b*
239 (E, F), *gabra3* (G, H), *gabra4* (I, J), and *gabra5* (K, L) were detected at this time point. Whole
240 mount lateral (A, C, E, G, I, K) views are shown along with dorsal views of the head (B, D, F, H,
241 J, L). The scale bar (A) is 0.1 mm. The brackets in G, I, and K indicates the regions shown at
242 higher magnification in the corresponding insets. Abbreviations: drc, dorsal rostral cluster; hbc,
243 hindbrain cluster; op, olfactory placode; SC, spinal cord; t, tegmentum; vcc, ventral caudal
244 cluster; vrc, ventral rostral cluster.

245

246 *gabra1*, *gabra3*, *gabra4*, and *gabra5* were each expressed in discrete cells in the brain
247 or spinal cord at 24 hpf. *gabra1* was detected in the olfactory placodes, the ventral rostral
248 cluster, the ventral caudal cluster and small clusters of cells within each rhombomere of the
249 hindbrain (Fig 2A, B). Although expressed in discrete cells, *gabra3* was widely expressed, and
250 found within the tegmentum, hindbrain, and ventral and intermediate domains in the spinal cord
251 (Fig 2G, H). *gabra4* was expressed in a diffuse manner throughout the brain but specifically
252 within a select population of ventral spinal cord cells (Fig 2I, J). The location and distribution of
253 these cells suggests that *gabra4* is expressed selectively within Kolmer-Agduhr (KA) neurons.
254 Lastly, *gabra5* is prominently expressed in small groups of cells in each rhombomere of the
255 hindbrain and in an intermediate domain of the spinal cord (Fig 2K, L). The location and
256 distribution of these spinal cord cells suggests that they are CoPA neurons.

257 The expression of all GABA_A receptor subunit genes, except for *gabra6a*, was observed
258 at 48 hpf (Fig 3). Similar to the pattern observed at 24 hpf, the *gabra2* paralogs were again
259 detected broadly, with little spatial restriction (Fig 3C-F). Similar to its expression at 24 hpf,
260 *gabra1* is prominently expressed in the olfactory bulbs, the pallium and discrete clusters of cells
261 in medial portions of the medulla (Fig 3A, B). *gabra3* was observed in relatively small clusters of
262 cells in the pallium, thalamus, and the medulla (Fig 3G, H). *gabra4* transcripts were identified in

263 distinct cells in the subpallium and lateral portions of the medulla. *gabra5* transcripts were
264 detected most prominently in the medulla. The location and expression this subunit suggests it
265 exhibits robust expression in the Mauthner cells, therefore $\alpha 5$ containing receptors are likely
266 some of the GABA_A isoforms identified previously via electrophysiology [25]. *gabra6b* is
267 expressed prominently in the olfactory bulbs.

268

269 **Fig 3. Expression of the GABA_A α subunits in the brain at 48hpf.** *gabra1* (A, B), *gabra2a* (C,
270 D), *gabra2b* (E, F) *gabra3* (G, H) *gabra4* (I, J) *gabra5* (K, L) and *gabra6b* (M, N) were detected
271 at this time point. Lateral (A, C, E, G, I, K, M) and dorsal (B, D, F, H, J, L, N) views are shown.
272 The scale bar (A) is 0.1 mm. Brackets in A, G, I, and K indicate the region shown in cross
273 sections within the insets. Abbreviations: cb, cerebellum; di, diencephalon; hth, hypothalamus;
274 M, Mauthner cell; mo, medulla oblongata; ob, olfactory bulb; p, pallium; sp, subpallium; t,
275 tegmentum; th, thalamus.

276

277 Although *gabra1*, *gabra3*, *gabra4*, and *gabra5* transcripts were all detected in the
278 medulla, they do not appear to be in the same cells. Instead, an intriguing medial to lateral
279 organization was observed. *gabra1* is expressed strongly in the most medial cells, *gabra3* and
280 *gabra5* transcripts were found in an intermediate domain, and *gabra4* was observed laterally
281 (compare insets in Fig A, G, I, K).

282 Transcripts for all α subunits are detected in larval zebrafish at 96 hpf (Fig 4). As with the
283 earlier developmental time points, *gabra2a* and *gabra2b* transcripts were detected broadly,
284 although *gabra2b* appears less diffuse and more discrete compared to at 24 and 48 hpf (Fig 4C-
285 F). *gabra1*, *gabra3*, and *gabra6b* are also expressed more broadly compared to earlier stages.
286 In contrast, *gabra4*, *gabra5* and *gabra6a* transcripts were detected in smaller groups of cells.

287 *gabra4* is expressed in a striped pattern in the outer nuclear layer of the retina, the posterior
288 tuberculum area, which is a portion of the diencephalon, and the tectum, cerebellum, and
289 medulla (Fig I, J). *gabra5* is expressed in discrete cells in the pallium, hypothalamus, cerebellum
290 and medulla (Fig K, L). Within the medulla, *gabra5* expression in the Mauthner Cells is robust,
291 as it is at earlier developmental stages.

292

293 **Fig 4. Expression of the GABA_A α subunits in the brain at 96hpf.** Transcripts encoding all α
294 subunits were detected at this time point. Lateral (A, C, E, G, I, K, M, O) and dorsal (B, D, F, H,
295 J, L, N, P) are shown The scale bar (A) is 0.1 mm. Abbreviations: cb, cerebellum; gcl, ganglion
296 cell layer; hth, hypothalamus; inl, inner nuclear layer; M, mauthner cell; mo, medulla oblongata;
297 ob, olfactory bulb; opn, optic nerve; p, pallium; pcl, photoreceptive cell layer; po, preoptic region;
298 poc, post optic commissure; pta, posterior tubercular area; sp, subpallium; t, tegmentum; te,
299 telencephalon; th, thalamus; to, tectum opticum;

300

301 Transcripts for several α subunits are expressed in the retina at 96 hpf, however most
302 are restricted to one or two cell layers (Fig 4). *gabra1* and *gabra6b* are both expressed in the
303 ganglion cell and outer nuclear layers, *gabra2a* and *gabra2b* both seem to be expressed
304 broadly, and *gabra3* and *gabra4* transcripts both are restricted to the outer nuclear layer.
305 *gabra6a* is expressed prominently in the photoreceptor cell layer, and does not demonstrate
306 robust expression outside of the retina.

307

308

309

310 DISCUSSION

311 In this study, we showed that the GABA_A receptor gene family exhibits a size and
312 diversity very similar to those found in mammals. We identified sequences for 23 GABA_A
313 receptor encoding genes, and phylogenetic analysis indicates that isoform are conserved
314 among vertebrates for the majority of subunits. We determined the expression of the eight
315 zebrafish α subunit encoding genes, the largest and most diverse gene family, and observed
316 that they are expressed in distinct, often overlapping expression patterns. Taken together, these
317 data argue that larval zebrafish can serve as a useful model to investigate the functional roles of
318 GABA_A receptor subtypes within developing neural circuits.

319 A previous study identified 23 GABA_A receptor genes in zebrafish. Our data largely
320 confirm their results, except ρ 3b, which is encoded by *gabrr3b*, was not reported. A α 3 and an
321 α 3-like gene were also described, but our analysis indicates that zebrafish likely contain only
322 one α 3 gene. Given the amount of zebrafish genome data that is available, it seems unlikely
323 that additional GABA_A receptor subunit encoding genes exist. However, similar to mammalian
324 systems, sequence data indicates that several zebrafish GABA_A receptor encoding genes are
325 alternatively spliced, which further enhances the already extensive diversity of receptor
326 isoforms.

327 The evolution of the GABA_A receptor gene family has been examined by using sequence
328 data from a wide variety of species, including humans, rodents, canary, chicken, frogs and
329 pufferfish, tunicates, *C. elegans* and *drosophila* [41, 42]. Our results are in line with these
330 studies in finding that mammalian θ and ϵ subunits are more distinct compared to other
331 subunits, such that their existence outside of the mammalian lineage is unclear [43]. Even
332 among mammalian species, for example comparing humans to rodents, these two subunits are
333 very diverse, which suggests they are evolving at a much faster rate than other subunits [44].

334 The roles of θ and ϵ subunits may be unique to mammals or, alternatively, fulfilled in zebrafish
335 and other species using other subtypes. Experiments to investigate the functional ability of
336 zebrafish subtypes to substitute for θ and ϵ in rodent systems could distinguish between these
337 possibilities.

338 The α subunits demonstrate a wide range of expression patterns across early zebrafish
339 development (Table 1). On one end of the spectrum are *gabra2a* and *gabra2b*, which appeared
340 to be expressed very broadly at each of the three time points examined. At the other end of the
341 spectrum is *gabra6a*, which we detected only at 96 hpf and mostly in the photoreceptor cell
342 layer of the retina. The other subunits fall between these extremes, with expression in discrete
343 cells in various brain and spinal cord regions. When GABA_A receptor α subtypes are expressed
344 in the same region, they often occupy different domains. For example, *gabra1*, *gabra3*, *gabra4*,
345 and *gabra5* are all expressed in the medulla, but are organized in medial to lateral stripes. The
346 zebrafish medulla has a structural and functional organization in which neurons of shared
347 neurotransmitter, phenotype, age, morphology and functional properties are arranged into
348 stripes [45, 46]. Another example of different GABA_A α subtypes occupying different domains
349 within the same region is found in the spinal cord. *gabra4* transcripts were found in ventral cells,
350 likely KA neurons, *gabra5*-expressing cells, likely CoPA neurons, were observed more dorsally,
351 and *gabra3* transcripts were detected more broadly in both ventral and more dorsal domains. It
352 is not yet clear how the expression of different GABA_A subtypes correspond to the
353 organizational stripes of the hindbrain or cell types in the spinal cord, but it seems likely that
354 some subtypes demonstrate cell-type specific expression and confer distinct responses to
355 GABA. Conclusive cell-type identification will require colabeling experiments that examine
356 GABA_A subunit expression along with specific markers or techniques that determine neuronal
357 identity by revealing cell morphology [47].

358

360 **Table 1** Shaded boxes indicate broad, diffuse expression was observed and dots indicate
361 punctate expression was detected.

362

363 The expression patterns of GABA_A α subunits are well-conserved across species. The
364 expression of five α subunits has been described in *Xenopus laevis* during development [48].
365 Consistent with our observations in zebrafish, $\alpha 2$ was found to be expressed broadly and early
366 in development. In fact, $\alpha 2$ was found to be deposited in eggs as a maternal message. $\alpha 6$
367 showed the latest onset of expression during development and demonstrated robust
368 photoreceptor expression. Likewise the expression patterns of $\alpha 1$, $\alpha 3$, and $\alpha 5$ are similar
369 between frogs and zebrafish. These similarities in expression extend to mammalian systems.
370 For example, in rat, similar to zebrafish $\alpha 2$, transcripts were detected relatively early in
371 development and were widespread, and $\alpha 6$ is expressed much later in development and
372 exhibits restricted expression [7]. One notable difference is that in mammals $\alpha 6$ transcripts were
373 detected in the cerebellum and retinal expression has not been reported, whereas in zebrafish
374 $\alpha 6a$ exhibits robust retinal expression and $\alpha 6b$ is more widespread. It is unclear how much
375 these differences in expression between zebrafish and mammals can be attributed to the
376 various development stages and tissues selected for investigation versus authentic differences
377 in patterns of expression.

378 Developing zebrafish are a powerful system for neural circuit analysis and, more
379 recently, they have been cultivated as a model of epilepsy [17, 49-53]. Although GABA_A
380 receptors are robust regulators of many neural circuits and play a central role in epilepsy
381 syndromes, GABA_A receptors have not been well characterized in zebrafish. By reporting the
382 diversity of zebrafish GABA_A receptor subunits and describing the expression patterns of the α
383 subfamily during early development, we have laid a foundation to leverage the strengths of the
384 zebrafish system to investigate isoform specific roles of GABA_A receptors and further develop

385 zebrafish as a model of epilepsy. In mammalian systems, genetic inactivation of individual α
386 subunits has revealed surprisingly subtle defects compared to what would be predicted from
387 pharmacological blockade. Compensatory upregulation of other subunits is thought to mask
388 some of the effects of inactivating a single subunit [54, 55]. The ability easily mutate multiple
389 genes in zebrafish may help overcome such compensatory changes in expression to shed new
390 light into the subtype specific roles of GABA_A receptors.

391 **ACKNOWLEDGEMENTS**

392 The authors thank members of the Downes lab, Kelly Anne McKeown at Westfield State
393 University, and Josef Trapani at Amherst College for helpful advice and discussion.

394 **REFERENCES**

- 395 1. Bowery NG, Smart TG. GABA and glycine as neurotransmitters: a brief history. *Br J*
396 *Pharmacol.* 2006;147 Suppl 1:S109-19. Epub 2006/01/13. doi: 10.1038/sj.bjp.0706443. PubMed
397 PMID: 16402094; PubMed Central PMCID: PMCPMC1760744.
- 398 2. Mohler H. GABA(A) receptor diversity and pharmacology. *Cell Tissue Res.*
399 2006;326(2):505-16. Epub 2006/08/29. doi: 10.1007/s00441-006-0284-3. PubMed PMID:
400 16937111.
- 401 3. Sigel E, Steinmann ME. Structure, function, and modulation of GABA(A) receptors. *J*
402 *Biol Chem.* 2012;287(48):40224-31. Epub 2012/10/06. doi: 10.1074/jbc.R112.386664. PubMed
403 PMID: 23038269; PubMed Central PMCID: PMCPMC3504738.
- 404 4. Chua HC, Chebib M. GABAA Receptors and the Diversity in their Structure and
405 Pharmacology. *Adv Pharmacol.* 2017;79:1-34. Epub 2017/05/23. doi:
406 10.1016/bs.apha.2017.03.003. PubMed PMID: 28528665.
- 407 5. Olsen RW, Sieghart W. International Union of Pharmacology. LXX. Subtypes of gamma-
408 aminobutyric acid(A) receptors: classification on the basis of subunit composition,
409 pharmacology, and function. Update. *Pharmacol Rev.* 2008;60(3):243-60. Epub 2008/09/16.
410 doi: 10.1124/pr.108.00505. PubMed PMID: 18790874; PubMed Central PMCID:
411 PMCPMC2847512.
- 412 6. Laurie DJ, Seeburg PH, Wisden W. The distribution of 13 GABAA receptor subunit
413 mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *J Neurosci.* 1992;12(3):1063-76.
414 Epub 1992/03/01. PubMed PMID: 1312132.

- 415 7. Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABAA receptor subunit
416 mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci*.
417 1992;12(11):4151-72. PubMed PMID: 1331359.
- 418 8. Wisden W, Laurie DJ, Monyer H, Seeburg PH. The distribution of 13 GABAA receptor
419 subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci*.
420 1992;12(3):1040-62. Epub 1992/03/01. PubMed PMID: 1312131.
- 421 9. Delgado-Lezama R, Loeza-Alcocer E, Andres C, Aguilar J, Guertin PA, Felix R.
422 Extrasynaptic GABA(A) receptors in the brainstem and spinal cord: structure and function. *Curr*
423 *Pharm Des*. 2013;19(24):4485-97. Epub 2013/01/31. PubMed PMID: 23360278.
- 424 10. Simon J, Wakimoto H, Fujita N, Lalande M, Barnard EA. Analysis of the set of GABA(A)
425 receptor genes in the human genome. *J Biol Chem*. 2004;279(40):41422-35. Epub 2004/07/20.
426 doi: 10.1074/jbc.M401354200. PubMed PMID: 15258161.
- 427 11. Olsen RW, Sieghart W. GABA A receptors: subtypes provide diversity of function and
428 pharmacology. *Neuropharmacology*. 2009;56(1):141-8. Epub 2008/09/02. doi:
429 10.1016/j.neuropharm.2008.07.045. PubMed PMID: 18760291; PubMed Central PMCID:
430 PMCPMC3525320.
- 431 12. Tretter V, Moss SJ. GABA(A) Receptor Dynamics and Constructing GABAergic
432 Synapses. *Front Mol Neurosci*. 2008;1:7. Epub 2008/10/24. doi: 10.3389/neuro.02.007.2008.
433 PubMed PMID: 18946540; PubMed Central PMCID: PMCPMC2526003.
- 434 13. Fritschy JM, Panzanelli P. GABAA receptors and plasticity of inhibitory
435 neurotransmission in the central nervous system. *Eur J Neurosci*. 2014;39(11):1845-65. Epub
436 2014/03/19. doi: 10.1111/ejn.12534. PubMed PMID: 24628861.
- 437 14. Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of
438 GABA(A) receptors. *Nat Rev Neurosci*. 2005;6(3):215-29. Epub 2005/03/02. doi:
439 10.1038/nrn1625. PubMed PMID: 15738957.
- 440 15. Egawa K, Fukuda A. Pathophysiological power of improper tonic GABA(A)
441 conductances in mature and immature models. *Front Neural Circuits*. 2013;7:170. Epub
442 2013/10/30. doi: 10.3389/fncir.2013.00170. PubMed PMID: 24167475; PubMed Central PMCID:
443 PMCPMC3807051.
- 444 16. Wolman M, Granato M. Behavioral genetics in larval zebrafish: learning from the young.
445 *Dev Neurobiol*. 2012;72(3):366-72. Epub 2012/02/14. doi: 10.1002/dneu.20872. PubMed PMID:
446 22328273.
- 447 17. McLean DL, Fetcho JR. Movement, technology and discovery in the zebrafish. *Curr Opin*
448 *Neurobiol*. 2011;21(1):110-5. Epub 2010/10/26. doi: 10.1016/j.conb.2010.09.011. PubMed
449 PMID: 20970321; PubMed Central PMCID: PMCPMC3057273.
- 450 18. Baraban SC, Taylor MR, Castro PA, Baier H. Pentylentetrazole induced changes in
451 zebrafish behavior, neural activity and c-fos expression. *Neuroscience*. 2005;131(3):759-68.
452 Epub 2005/02/26. doi: 10.1016/j.neuroscience.2004.11.031. PubMed PMID: 15730879.
- 453 19. Stehr CM, Linbo TL, Incardona JP, Scholz NL. The developmental neurotoxicity of
454 fipronil: notochord degeneration and locomotor defects in zebrafish embryos and larvae. *Toxicol*
455 *Sci*. 2006;92(1):270-8. Epub 2006/04/14. doi: 10.1093/toxsci/kfj185. PubMed PMID: 16611622.
- 456 20. Connaughton VP, Nelson R, Bender AM. Electrophysiological evidence of GABAA and
457 GABAC receptors on zebrafish retinal bipolar cells. *Vis Neurosci*. 2008;25(2):139-53. Epub
458 2008/04/30. doi: 10.1017/S0952523808080322. PubMed PMID: 18442437; PubMed Central
459 PMCID: PMCPMC4272819.

- 460 21. Del Bene F, Wyart C, Robles E, Tran A, Looger L, Scott EK, et al. Filtering of visual
461 information in the tectum by an identified neural circuit. *Science*. 2010;330(6004):669-73. Epub
462 2010/10/30. doi: 10.1126/science.1192949. PubMed PMID: 21030657; PubMed Central PMCID:
463 PMCPMC3243732.
- 464 22. Mack-Bucher JA, Li J, Friedrich RW. Early functional development of interneurons in the
465 zebrafish olfactory bulb. *Eur J Neurosci*. 2007;25(2):460-70. Epub 2007/02/08. doi:
466 10.1111/j.1460-9568.2006.05290.x. PubMed PMID: 17284187.
- 467 23. Hubbard JM, Bohm UL, Prendergast A, Tseng PB, Newman M, Stokes C, et al.
468 Intraspinal Sensory Neurons Provide Powerful Inhibition to Motor Circuits Ensuring Postural
469 Control during Locomotion. *Curr Biol*. 2016;26(21):2841-53. Epub 2016/10/11. doi:
470 10.1016/j.cub.2016.08.026. PubMed PMID: 27720623.
- 471 24. Triller A, Rostaing P, Korn H, Legendre P. Morphofunctional evidence for mature
472 synaptic contacts on the Mauthner cell of 52-hour-old zebrafish larvae. *Neuroscience*.
473 1997;80(1):133-45. Epub 1997/09/01. PubMed PMID: 9252227.
- 474 25. Roy B, Ali DW. Multiple types of GABAA responses identified from zebrafish Mauthner
475 cells. *Neuroreport*. 2014;25(15):1232-6. Epub 2014/08/28. doi:
476 10.1097/WNR.0000000000000258. PubMed PMID: 25162782.
- 477 26. Cocco A, Ronnberg AM, Jin Z, Andre GI, Vossen LE, Bhandage AK, et al.
478 Characterization of the gamma-aminobutyric acid signaling system in the zebrafish (*Danio rerio*
479 Hamilton) central nervous system by reverse transcription-quantitative polymerase chain
480 reaction. *Neuroscience*. 2017;343:300-21. Epub 2016/07/28. doi:
481 10.1016/j.neuroscience.2016.07.018. PubMed PMID: 27453477.
- 482 27. Hortopan GA, Dinday MT, Baraban SC. Spontaneous seizures and altered gene
483 expression in GABA signaling pathways in a mind bomb mutant zebrafish. *J Neurosci*.
484 2010;30(41):13718-28. Epub 2010/10/15. doi: 10.1523/JNEUROSCI.1887-10.2010. PubMed
485 PMID: 20943912; PubMed Central PMCID: PMCPMC2962868.
- 486 28. Gonzalez-Nunez V. Role of *gabra2*, GABAA receptor alpha-2 subunit, in CNS
487 development. *Biochem Biophys Reports*. 2015;3:11.
- 488 29. Baxendale S, Holdsworth CJ, Meza Santoscoy PL, Harrison MR, Fox J, Parkin CA, et al.
489 Identification of compounds with anti-convulsant properties in a zebrafish model of epileptic
490 seizures. *Dis Model Mech*. 2012;5(6):773-84. Epub 2012/06/26. doi: 10.1242/dmm.010090.
491 PubMed PMID: 22730455; PubMed Central PMCID: PMCPMC3484860.
- 492 30. Thisse B, Thisse C. . Fast Release Clones: A High Throughput Expression Analysis.
493 ZFIN Direct Data Submission. 2004.
- 494 31. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of
495 progressive multiple sequence alignment through sequence weighting, position-specific gap
496 penalties and weight matrix choice. *Nucleic Acids Res*. 1994;22(22):4673-80. Epub 1994/11/11.
497 PubMed PMID: 7984417; PubMed Central PMCID: PMCPMC308517.
- 498 32. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
499 and methods to estimate maximum-likelihood phylogenies: assessing the performance of
500 PhyML 3.0. *Syst Biol*. 2010;59(3):307-21. Epub 2010/06/09. doi: 10.1093/sysbio/syq010.
501 PubMed PMID: 20525638.
- 502 33. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic
503 development of the zebrafish. *Dev Dyn*. 1995;203(3):253-310. PubMed PMID: 8589427.

- 504 34. Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount zebrafish
505 embryos. *Nat Protoc.* 2008;3(1):59-69. PubMed PMID: 18193022.
- 506 35. Mueller T, Wullimann MF. Atlas of early zebrafish brain development : a tool for
507 molecular neurogenetics. Second edition. ed. xiii, 244 pages p.
- 508 36. Folgueira M, Bayley P, Navratilova P, Becker TS, Wilson SW, Clarke JD.
509 Morphogenesis underlying the development of the everted teleost telencephalon. *Neural Dev.*
510 2012;7:32. Epub 2012/09/20. doi: 10.1186/1749-8104-7-32. PubMed PMID: 22989074; PubMed
511 Central PMCID: PMC3520737.
- 512 37. Sabaliauskas NA, Foutz CA, Mest JR, Budgeon LR, Sidor AT, Gershenson JA, et al.
513 High-throughput zebrafish histology. *Methods.* 2006;39(3):246-54. Epub 2006/07/28. doi:
514 10.1016/j.ymeth.2006.03.001. PubMed PMID: 16870470.
- 515 38. Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, et al. Zebrafish hox clusters
516 and vertebrate genome evolution. *Science.* 1998;282(5394):1711-4. Epub 1998/11/30. PubMed
517 PMID: 9831563.
- 518 39. Postlethwait JH, Yan YL, Gates MA, Horne S, Amores A, Brownlie A, et al. Vertebrate
519 genome evolution and the zebrafish gene map. *Nat Genet.* 1998;18(4):345-9. Epub 1998/04/16.
520 doi: 10.1038/ng0498-345. PubMed PMID: 9537416.
- 521 40. Saint-Amant L, Drapeau P. Time course of the development of motor behaviors in the
522 zebrafish embryo. *J Neurobiol.* 1998;37(4):622-32. PubMed PMID: 9858263.
- 523 41. Tsang SY, Ng SK, Xu Z, Xue H. The evolution of GABAA receptor-like genes. *Mol Biol*
524 *Evol.* 2007;24(2):599-610. Epub 2006/12/01. doi: 10.1093/molbev/msl188. PubMed PMID:
525 17135332.
- 526 42. Darlison MG, Pahal I, Thode C. Consequences of the evolution of the GABA(A) receptor
527 gene family. *Cell Mol Neurobiol.* 2005;25(3-4):607-24. Epub 2005/08/03. doi: 10.1007/s10571-
528 005-4004-4. PubMed PMID: 16075381.
- 529 43. Martyniuk CJ, Aris-Brosou S, Drouin G, Cahn J, Trudeau VL. Early evolution of
530 ionotropic GABA receptors and selective regimes acting on the mammalian-specific theta and
531 epsilon subunits. *PLoS One.* 2007;2(9):e894. Epub 2007/09/20. doi:
532 10.1371/journal.pone.0000894. PubMed PMID: 17878929; PubMed Central PMCID:
533 PMCPMC1975676.
- 534 44. Sinkkonen ST, Hanna MC, Kirkness EF, Korpi ER. GABA(A) receptor epsilon and theta
535 subunits display unusual structural variation between species and are enriched in the rat locus
536 ceruleus. *J Neurosci.* 2000;20(10):3588-95. Epub 2000/05/11. PubMed PMID: 10804200.
- 537 45. Higashijima S, Mandel G, Fetcho JR. Distribution of prospective glutamatergic,
538 glycinergic, and GABAergic neurons in embryonic and larval zebrafish. *J Comp Neurol.*
539 2004;480(1):1-18. Epub 2004/10/30. doi: 10.1002/cne.20278. PubMed PMID: 15515020.
- 540 46. Kinkhabwala A, Riley M, Koyama M, Monen J, Satou C, Kimura Y, et al. A structural and
541 functional ground plan for neurons in the hindbrain of zebrafish. *Proc Natl Acad Sci U S A.*
542 2011;108(3):1164-9. Epub 2011/01/05. doi: 10.1073/pnas.1012185108. PubMed PMID:
543 21199947; PubMed Central PMCID: PMCPMC3024665.
- 544 47. Downes GB, Waterbury JA, Granato M. Rapid in vivo labeling of identified zebrafish
545 neurons. *Genesis.* 2002;34(3):196-202. PubMed PMID: 12395384.
- 546 48. Kaeser GE, Rabe BA, Saha MS. Cloning and characterization of GABAA alpha subunits
547 and GABAB subunits in *Xenopus laevis* during development. *Dev Dyn.* 2011;240(4):862-73.

- 548 Epub 2011/03/09. doi: 10.1002/dvdy.22580. PubMed PMID: 21384470; PubMed Central
549 PMCID: PMCPMC3071254.
- 550 49. Fetcho JR, Higashijima S, McLean DL. Zebrafish and motor control over the last decade.
551 Brain Res Rev. 2008;57(1):86-93. Epub 2007/09/11. doi: 10.1016/j.brainresrev.2007.06.018.
552 PubMed PMID: 17825423; PubMed Central PMCID: PMCPMC2237884.
- 553 50. Friedrich RW, Genoud C, Wanner AA. Analyzing the structure and function of neuronal
554 circuits in zebrafish. Front Neural Circuits. 2013;7:71. Epub 2013/05/01. doi:
555 10.3389/fncir.2013.00071. PubMed PMID: 23630467; PubMed Central PMCID:
556 PMCPMC3632777.
- 557 51. Hale ME, Katz HR, Peek MY, Fremont RT. Neural circuits that drive startle behavior,
558 with a focus on the Mauthner cells and spiral fiber neurons of fishes. J Neurogenet.
559 2016;30(2):89-100. Epub 2016/06/16. doi: 10.1080/01677063.2016.1182526. PubMed PMID:
560 27302612.
- 561 52. Griffin A, Krasniak C, Baraban SC. Advancing epilepsy treatment through personalized
562 genetic zebrafish models. Prog Brain Res. 2016;226:195-207. Epub 2016/06/22. doi:
563 10.1016/bs.pbr.2016.03.012. PubMed PMID: 27323944.
- 564 53. Cunliffe VT. Building a zebrafish toolkit for investigating the pathobiology of epilepsy and
565 identifying new treatments for epileptic seizures. J Neurosci Methods. 2016;260:91-5. Epub
566 2015/07/30. doi: 10.1016/j.jneumeth.2015.07.015. PubMed PMID: 26219659.
- 567 54. Kralic JE, Sidler C, Parpan F, Homanics GE, Morrow AL, Fritschy JM. Compensatory
568 alteration of inhibitory synaptic circuits in cerebellum and thalamus of gamma-aminobutyric acid
569 type A receptor alpha1 subunit knockout mice. J Comp Neurol. 2006;495(4):408-21. Epub
570 2006/02/18. doi: 10.1002/cne.20866. PubMed PMID: 16485284.
- 571 55. Peng Z, Zhang N, Chandra D, Homanics GE, Olsen RW, Houser CR. Altered
572 localization of the delta subunit of the GABAA receptor in the thalamus of alpha4 subunit
573 knockout mice. Neurochem Res. 2014;39(6):1104-17. Epub 2013/12/20. doi: 10.1007/s11064-
574 013-1202-1. PubMed PMID: 24352815; PubMed Central PMCID: PMCPMC4024081.

575

576

577 **Supporting Information**

578 **S1 Fig. RT-PCR results suggest that zebrafish contain only one $\alpha 3$ gene**

579 (A) Schematic of a *gabra3* cDNA. The portions that are identical to *gabra3*-like and *gabra3*
580 sequences are shaded green. The white region that connects the two was identified through RT-
581 PCR. The location of the three primers used for PCR are shown as numbered arrows. (B) RT-PCR
582 results. Primers 2 and 3 served as a positive control since they amplify sequence from the known
583 *gabra3* region. Primers 1 and 3 amplify a previously unknown region that links *gabra 3*-like and
584 *gabra3* sequence, showing that they are contained within same transcript and likely portions of the
585 same gene.

586

587 **S1 Table. GABA_A α subunit antisense RNA probe information.** Note- Sizes are shown in base
588 pairs. The probe start site is defined here as the start codon of the open reading frame. Negative
589 values indicate that the probe contains 5' untranslated sequence. Some probes were generated via
590 RT-PCR with a primer that contains a RNA polymerase promoter and subcloning into vectors wasn't
591 performed.

592







