

TagMap  
Tn5 Tagmentation-based pBac mapping  
David L. Stern  
January 11, 2016

## Protocol - Preliminaries

### **1 - Anneal the oligonucleotides**

Combine:

Rx1 (adapter 1 10 uM)

10uL Tn5ME-A (100 uM)  
10uL Tn5MErev-InT (100 uM)  
80uL Reassociation Buffer

Mix well.

Anneal primers in a thermal cycler with the following Reassociation Program

Step	Temp	Time
1	95°C	10 min
2	90°C	1 min
3	Reduce temp by 1°C/cycle 60 times	
4	4°C	Hold

### **2 - Dilute Tn5 to 20ng/uL**

e.g.

1 uL Tn5 protein (200 ng / uL)  
9 uL Reassociation buffer

Mix well by pipetting 10 times.

### **3 - Dilute the adaptors.**

1 uL Adaptor 1 (10 uM)  
9 uL H2O

Mix by pipetting 10 times.

#### **4 - Pre-charge the Tn5**

Combine, in the following order, mix after each addition:

21 uL Tn5 (20 ng / uL)

10 uL Glycerol

10 uL Diluted Adaptors (1uM)

Pre-charge at 37°C for 30 minutes

#### **5 - Tagment**

Combine:

1 uL Precharged Tn5

1 uL DNA (10 ng / uL *or less!*)

2 uL 5 X TAPS

6 uL H<sub>2</sub>O

Mix well by pipetting up and down 10 X.

Incubate at 55°C for 7 minutes.

#### **6 - Kill the Tn5**

Add to each reaction:

2.5 uL 0.2% SDS

Incubate at 55° 7 min

## 7 - PCR 1

1uL	Tagmentation reaction
1uL	5uM Tn5ME-B-pBac-142R
1uL	5uM Tn5ME-B-pBac-3287F
1uL	5uM A_idx_i5 primer
1uL	4 mM dNTP
4uL	5X Phusion Reaction Buffer
0.5uL	Phusion Polymerase
10.5uL	H <sub>2</sub> O

### Thermocycler settings

Step	Temp	Time
1	95°C	5 min
1	95°C	15 sec
2	60°C	15 sec
3	72°C	1 min
4	Cycle to step 3 19 times	
5	72°C	2 min
6	4°C	Hold

## 7.5 — Optional Ampure cleanup of pooled reactions

Pool 5uL of each reaction = 480uL

100 uL	Pooled reactions
80 uL	Ampure beads

Bind, wash 80% Ethanol X 2  
Resuspend in 100 uL H<sub>2</sub>O

## 8 - PCR 2

1 uL	PCR1 Reaction
1 uL	10 uM Primer 1 - Tn5-Illumina-Primer1 (FC2)
1 uL	10 uM Primer 2 - i7_idx-primer
1 uL	dNTPs
4 uL	Phusion 5X Buffer
0.5uL	Phusion Polymerase
11.5uL	H <sub>2</sub> O

Cycle 20 times with same thermocycler settings

## **12 – Clean up product - Ampure**

Pool 10uL of each reaction

0.8 : 1          Ampure : pooled reactions

Wash 2 X 80% Ethanol

Resuspend in H<sub>2</sub>O volume of original pooled reactions

## **13 – Quantify final library**

Reassociation Buffer – *store at R.T.*

10 mM Tris pH 8.0

50 mM NaCl

1 mM EDTA

5x TAPS-DMF buffer from Picelli paper

50 mM TAPS-NaOH,

25 mM MgCl<sub>2</sub>,

50% v/v DMF (pH 8.5) at 25°C

-100 g TAPS in 500 ml H<sub>2</sub>O, pH = 9.9 -Add 5-10 ml concentrate HCl to get to pH 8.5

-complete to 754 ml for 500mM (Paper says TAPS-NaOH but I added HCl instead)