

Supplementary Information for:

Absolute quantification of translational regulation and burden using combined sequencing approaches

Thomas E. Gorochowski, Irina Chelysheva, Mette Eriksen, Priyanka Nair, Steen Pedersen, Zoya Ignatova

Supplementary Figures	Page
Figure S1: Sequencing data exhibit a good correlation between biological replicates	2
Figure S2: Comparison of ribosome protected fragments (RPFs) mapping to first and second half of each coding region	3
Supplementary Tables	
Table S1: Measurements of cellular parameters	4
Table S2: Sequencing and read mapping statistics	5




Figure S1: Sequencing data exhibit a good correlation between biological replicates. Correlation of the RNA-seq and Ribo-seq data of two biological replicates from induced and non-induced cells expressing either LacZ or LacZ-PK. R^2 , Pearson correlation coefficient.




Figure S2: Comparison of ribosome protected fragments (RPFs) mapping to first and second half of each coding region. Each point represents a coding region. First 51 bp and last 9 bp not included in the calculation. R², Pearson correlation coefficient.

Table S1: Measurements of cellular parameters

Measurement	Start of induction		End of induction ^a	
	LacZ	PK-LacZ	LacZ	PK-LacZ
Doubling time (minutes)	85	85	– ^b	– ^b
Cell count	1.6×10^9	1.6×10^9	1.6×10^9	1.6×10^9
Protein mass (grams/cell)	4.13×10^{-13}	4.10×10^{-13}	4.70×10^{-13}	4.30×10^{-13}
Mass of LacZ/PK-LacZ as total of cell	–	–	14%	5.8%

a. Induction of constructs lasted 10 min for LacZ and 15 min for PK-LacZ before samples were taken.

b. No measurable growth during the induction period.

Table S2: Sequencing and read mapping statistics

Sample ^a				Total # of reads ^b	Multi-mapped reads ^c	Non-coding ^c	Uniquely mapped to mRNA ^c
PK-LacZ	RNA-seq	non-induced	R#1	66066771	29796556 (45.10%)	2876188 (4.35%)	33394027 (50.55%)
PK-LacZ	RNA-seq	non-induced	R#2	39198111	18025740 (45.99%)	1673252 (4.27%)	19499119 (49.75%)
PK-LacZ	RNA-seq	induced	R#1	70004102	28545341 (40.78%)	1982506 (2.83%)	39476255 (56.39%)
PK-LacZ	RNA-seq	induced	R#2	68146152	27782049 (40.77%)	1931069 (2.83%)	38433034 (56.40%)
PK-LacZ	Ribo-seq	non-induced	R#1	59384989	25736660 (43.34%)	3034480 (5.11%)	30613849 (51.55%)
PK-LacZ	Ribo-seq	non-induced	R#2	62798001	23567687 (37.53%)	6009235 (9.57%)	33221079 (52.90%)
PK-LacZ	Ribo-seq	induced	R#1	56568122	27566889 (48.73%)	2387216 (4.22%)	26614017 (47.05%)
PK-LacZ	Ribo-seq	induced	R#2	54581989	20143633 (36.91%)	3541469 (6.49%)	30896887 (56.61%)
LacZ	RNA-seq	non-induced	R#1	192586734	90755510 (47.12%)	7758247 (4.03%)	94072977 (48.85%)
LacZ	RNA-seq	non-induced	R#2	53735584	28463851 (52.97%)	3382171 (6.29%)	21889562 (40.74%)
LacZ	RNA-seq	induced	R#1	190867334	91396079 (47.88%)	6916496 (3.62%)	92554759 (48.49%)
LacZ	RNA-seq	induced	R#2	52249590	25660126 (49.11%)	2776669 (5.31%)	23812795 (45.58%)
LacZ	Ribo-seq	non-induced	R#1	183013690	124093781 (67.81%)	11347972 (6.20%)	47571937 (25.99%)
LacZ	Ribo-seq	non-induced	R#2	53330842	25579314 (47.96%)	4072967 (7.64%)	23678561 (44.40%)
LacZ	Ribo-seq	induced	R#1	199707963	120352995 (60.26%)	7716217 (3.86%)	71638751 (35.87%)
LacZ	Ribo-seq	induced	R#2	41870343	31443591 (75.10%)	1843808 (4.40%)	8582944 (20.50%)

a. Biological replicates are denoted “R#1” and “R#2” and “induced” relates to presence of IPTG (1 mM).

b. Reads aligning to more than one sequence including tRNA and rRNA were excluded from the data.

c. Sequencing reads were mapped to the genome of *E. coli* K-12 MG1655 strain.