

TagMap
Tn5 Tagmentation-based pBac mapping
David L. Stern
February 2, 2016

Protocol - Preliminaries

1 - Anneal the oligonucleotides

Combine:

Rx1 (adapter 1 10 uM)

10uL Tn5ME-A (100 uM)
10uL Tn5MErev-InT (100 uM)
80uL Reassociation Buffer

Mix well.

Anneal primers in a thermal cycler with the following Reassociation Program

Step	Temp	Time
1	95°C	10 min
2	90°C	1 min
3	Reduce temp by 1°C/cycle 60 times	
4	4°C	Hold

2 - Dilute Tn5 to 20ng/uL

e.g.
1 uL Tn5 protein (200 ng / uL)
9 uL Reassociation buffer

Mix well by pipetting 10 times.

3 - Dilute the adaptors.

1 uL Adaptor 1 (10 uM)
9 uL H2O

Mix by pipetting 10 times.

4 - Pre-charge the Tn5

Combine, in the following order, mix after each addition:

21 uL Tn5 (20 ng / uL)

10 uL Glycerol

10 uL Diluted Adaptors (1uM)

Pre-charge at 37°C for 30 minutes

5 - Tagment

Combine:

1 uL Precharged Tn5

1 uL DNA (10 ng / uL *or less!*)

2 uL 5 X TAPS

6 uL H₂O

Mix well by pipetting up and down 10 X.

Incubate at 55°C for 7 minutes.

6 - Kill the Tn5

Add to each reaction:

2.5 uL 0.2% SDS

Incubate at 55° 7 min

7 - PCR 1

1uL Tagmentation reaction
1uL 5uM Tn5ME-B-pBac-142R
1uL 5uM Tn5ME-B-pBac-3287F
1uL 5uM A_idx_i5 primer
1uL 4 mM dNTP
4uL 5X Phusion Reaction Buffer
0.5uL Phusion Polymerase
10.5uL H₂O

Thermocycler settings

Step	Temp	Time
1	95°C	5 min
1	95°C	15 sec
2	60°C	15 sec
3	72°C	1 min
4	Cycle to step 3 19 times	
5	72°C	2 min
6	4°C	Hold

8 - PCR 2

1 uL PCR1 Reaction
1 uL 10 uM Primer 1 - Tn5-Illumina-Primer1 (FC2)
1 uL 10 uM Primer 2 - i7_idx-primer
1 uL dNTPs
4 uL Phusion 5X Buffer
0.5uL Phusion Polymerase
11.5uL H₂O

Cycle 15 times with same thermocycler settings as PCR 1

12 - Clean up product - Ampure

Pool 10uL of each reaction

0.8 : 1 Ampure : pooled reactions

Wash 2 X 80% Ethanol

Resuspend in H₂O volume of original pooled reactions

13 - Quantify final library

Reassociation Buffer – *store at R.T.*

10 mM Tris pH 8.0

50 mM NaCl

1 mM EDTA

5x TAPS-DMF buffer from Picelli paper

50 mM TAPS-NaOH,

25 mM MgCl₂,

50% v/v DMF (pH 8.5) at 25°C

-100 g TAPS in 500 ml H₂O, pH = 9.9 -Add 5-10 ml concentrate HCl to get to pH 8.5

-complete to 754 ml for 500mM (Paper says TAPS-NaOH but I added HCl instead)