




Irinotecan




Mechlorethamine



Paraquat










Figure S1 Dose-response High Throughput Assay. (A) Raw phenotypic values for four divergent strains, CB4856 (blue), DL238 (green), JU258 (pink), and N2 (orange), in response to various concentrations of each toxin are plotted as Tukey box plots. Toxin responses for traits representing three distinct population parameters- brood size (norm.n), mean animal length (mean.TOF), and mean optical density (mean.norm.EXT) are shown. The x-axis indicates either the control (water or DMSO) or the concentration (μM) of the toxin, and the y-axis shows the raw phenotypic value for each trait.


Cadmium




Carmustine




Chlorothalonil




Chlorpyrifos




Cisplatin




Copper




Diquat




Fluoxetine




FUDR




Irinotecan




Mechlorethamine




Paraquat




Silver



Topotecan



Tunicamycin



Vincristine











Figure S2 Trait correlations with RIAIL phenotypes. Correlation coefficients (r^2) between all pairwise combinations of traits based on the RIAIL phenotypes for the indicated concentration of the toxin are plotted as a heat map. Traits are labeled on the x- and y-axes, organized by hierarchical clustering (top) of the correlation matrix (bottom) for each toxin. Colors represent correlation coefficients increasing from green to yellow to pink to white.



A cadmium.PC2**B****C**

A cadmium.PC3**B****C**


V:18354580








A carmustine.PC1**B****C**




V:11869708








A carmustine.PC2**B****C**

I:5428703




A carmustine.PC3**B****C**


A carmustine.PC4**B****C**

A carmustine.PC6**B****C**


IV:6077574




A chlorothalonil.PC1




B




C




A chlorothalonil.PC2




B




C




A chlorothalonil.PC3




B




C



A chlorothalonil.PC4




B




C


V:19897614



A chlorothalonil.PC5




B




C


X:14089716




A chlorpyrifos.PC1




B




C



A chlorpyrifos.PC2




B




C


IV:15568713




A chlorpyrifos.PC5




B




C



A chlorpyrifos.PC6





B




C

V:16837198







A cisplatin.PC3




B




C



A cisplatin.PC4




B




C


V:14642834




A copper.PC2






B




C




A copper.PC4**B****C**


A copper.PC6






B




C




A diquat.PC2**B****C**

A diquat.PC5




B




C

X:12750794



A fluoxetine.PC1




C


IV:16442619

IV:4526752


V:3819945



A fluoxetine.PC5






B




C


IV:16259338






A FUdR.PC3**B****C**


IV:9715431






A irinotecan.PC2**B****C**

A irinotecan.PC3







B




C

V:4500931




A irinotecan.PC5**B****C**


A mechlorethamine.PC2





B




C




A mechlorethamine.PC7


**B****C**

IV:2052489







A silver.PC1









B





C




A silver.PC3**B****C**


A silver.PC4**B****C**

A silver.PC5**B****C**


IV:4009430




A topotecan.PC1






B




C




A topotecan.PC2**B****C**


A tunicamycin.PC1




B




C




A tunicamycin.PC3




B




C



A tunicamycin.PC4






B




C




V:4296845



A vincristine.PC1**B****C**

I:10163992



A vincristine.PC5**B****C**








Figure S3 Linkage mapping results and variance component contributions for 47 principal components with a significant QTL. **(A)** The maximum LOD score (y-axis) for each marker across the genome (x-axis), split by chromosome, is plotted as a black line. Significant QTL are indicated by a red triangle at the marker with the maximum LOD score and a light blue ribbon spanning the 95% confidence interval. The estimated phenotypic variation in the RIAILs explained by each QTL is indicated above each peak. **(B)** The proportion of phenotypic variation across the RIAILs caused by additive (light blue) and interactive (dark blue) genetic components are indicated. Solid and dashed error bars show the standard error around additive and interactive component estimates, respectively. **(C)** For each QTL plotted in A, the residual phenotypes (y-axis) of RIAILs split by genotype at the marker with the maximum LOD score (x-axis) are plotted as Tukey box plots. Data from RIAILs with the N2 or the CB4856 allele at the QTL peak marker are colored in orange and blue, respectively.



carmustine.PC1









carmustine.PC6


NIL – No parental diff.




NIL – Unidirectional





chlorothalonil.PC1






chlorothalonil.PC2


NIL – No parental diff.




chlorothalonil.PC3




NIL – Bidirectional




NIL – Unidirectional




NIL – Unidirectional




NIL – No parental diff.




NIL – Miscellaneous




cisplatin.PC1




NIL – No parental diff.




NIL – Unidirectional




NIL – Unidirectional




NIL – Unidirectional




NIL – Unidirectional





NIL – No parental diff.




CSS – No parental diff.









cisplatin.PC3




cisplatin.PC4




fluoxetine.PC1




NIL – No parental diff.




NIL – Miscellaneous




NIL – No parental diff.




fluoxetine.PC5




NIL – No parental diff.




NIL – Miscellaneous




NIL – No parental diff.





irinotecan.PC2




paraquat.PC1













silver.PC1









silver.PC3




NIL – Miscellaneous




NIL – Unidirectional




NIL – No effect




silver.PC4




silver.PC5




tunicamycin.PC1




NIL – Unidirectional




NIL – Bidirectional




NIL – Unidirectional




NIL – No parental diff.




NIL – Bidirectional




NIL – Unidirectional




NIL – Bidirectional




NIL – Unidirectional




NIL – Unidirectional



NIL – No parental diff.



NIL – Unidirectional



tunicamycin.PC3





Figure S5 NIL and CSS phenotypes. For traits in which NILs and CSSs were assayed, the residual phenotypic values (y-axis) of N2, CB4856, and reciprocal introgressed NILs or CSSs are plotted as Tukey box plots. Different letters represent significant differences ($p < 0.05$) between strains (TukeyHSD). The genotype of each strain on the x-axis is modeled by the colored rectangles beneath the plots (N2 genotypes are orange, CB4856 genotypes are blue). The category assigned to each test of recapitulation of QTL effect is marked above each NIL/CSS phenotype plot. Traits are grouped by the principal component they were selected to represent, based on correlation clusters from Figure S2.



Category	CSS	NIL
Recapitulation	Recapitulation	Recapitulation
No parental effect	No parental difference	No parental difference
No QTL effect	No QTL effect	No QTL effect
Inter-chromosomal external	Uni-directional or bi-directional transgressive phenotype	Recapitulation or no QTL effect
Inter-chromosomal internal	Uni-directional or bi-directional transgressive phenotype	Uni-directional or bi-directional transgressive phenotype
Intra-chromosomal	Recapitulation or no QTL effect	Uni-directional or bi-directional transgressive phenotype

Figure S6 NIL and CSS trait categorizations. (A) Flowchart for categorizing traits from the NIL or CSS tests of recapitulation of QTL effects. (B) Six potential categories for chromosome V traits tested in both NIL and CSS assays with a significant and consistent parental phenotypic split across both assays. The miscellaneous category is not depicted, but encompasses any other combination of NIL and CSS assay results.




Figure S7 Categorization of correlated traits. The y-axis shows the principal component tested in NIL assays, split by hotspot. Each color represents the NIL assay categorization for each trait within a correlation cluster - either recapitulation (green), no QTL effect (red), unidirectional transgressive phenotype (light blue), bidirectional transgressive phenotype (dark blue), miscellaneous (light grey) or no significant parental difference (dark grey). The percent of all traits within the correlation cluster for each principal component that falls within a given category is shown on the y-axis. Numbers on the right of each bar indicate the number of traits within a correlation cluster.